

**This Page Is Inserted by IFW Operations  
and is not a part of the Official Record**

## **BEST AVAILABLE IMAGES**

**Defective images within this document are accurate representations of the original documents submitted by the applicant.**

**Defects in the images may include (but are not limited to):**

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**





## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification<sup>7</sup> :</b> <b>A61K 48/00, C12N 15/63, 15/00, C07H 21/04</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/23116</b> <b>(43) International Publication Date:</b> 27 April 2000 (27.04.00)
<b>(21) International Application Number:</b> PCT/US99/24495 <b>(22) International Filing Date:</b> 19 October 1999 (19.10.99)  <b>(30) Priority Data:</b> 60/104,994           20 October 1998 (20.10.98)       US 60/125,974           24 March 1999 (24.03.99)       US 09/364,862           30 July 1999 (30.07.99)       US  <b>(71) Applicant (for all designated States except US):</b> AVIGEN, INC. [US/US]; #1000, 1201 Harbor Bay Parkway, Alameda, CA 94502 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> COUTO, Linda, B. [US/US]; 7814 Oak Creek Drive, Pleasanton, CA 94588 (US). COLOSI, Peter, C. [US/US]; 2016 Encinal Avenue, Alameda, CA 94501 (US).  <b>(74) Agents:</b> MACKNIGHT, Kamrin, T. et al.; Medlen & Carroll, LLP, Suite 2200, 220 Montgomery Street, San Francisco, CA 94104 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> ADENO-ASSOCIATED VECTORS FOR EXPRESSION OF FACTOR VIII BY TARGET CELLS  <b>(57) Abstract</b> <p>The present invention provides improved viral vectors useful for the expression of genes at high levels in human cells. In particular, the present invention provides adeno-associated vectors (AAV) suitable for gene therapy. These vectors are capable of delivering nucleic acid containing constructs which result in the production of full-length therapeutic levels of biologically active Factor VIII in the recipient individual <i>in vivo</i>. The present invention also provides pharmaceutical compositions comprising such AAV vectors, as well as methods for making and using these constructs.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## ADENO-ASSOCIATED VECTORS FOR EXPRESSION OF FACTOR VIII BY TARGET CELLS

This application claims priority benefit of U.S. provisional application Nos.  
5 60/125,974 filed March 24, 1999 and 60/104,994 filed October 20, 1998, which are  
hereby incorporated by reference in their entireties.

### FIELD OF THE INVENTION

The present invention relates to AAV vectors suitable for hemophilia gene  
10 therapy. More particularly, these AAV vectors are suitable for delivering nucleic acids  
encoding Factor VIII into a recipient subject suffering from hemophilia A, such that  
the subject's blood is able to clot.

### BACKGROUND

15 Hemophilia is a genetic disease characterized by a blood clotting deficiency. In  
hemophilia A (classic hemophilia, Factor VIII deficiency), an X-chromosome-linked  
genetic defect disrupts the gene encoding Factor VIII, a plasma glycoprotein, which is  
a key component in the blood clotting cascade. Human Factor VIII is synthesized as a  
single chain polypeptide, with a predicted molecular weight of 265 kDa. The Factor  
20 VIII gene codes for 2351 amino acids, and the protein has six domains, designated  
from the amino to the carboxy terminus as A1-A2-B-A3-C1-C2 (Wood *et al.*, Nature  
312:330 [1984]; Vehar *et al.*, Nature 312:337 [1984]; and Toole *et al.*, Nature 312:342  
[1984]). Human Factor VIII is processed within the cell to yield a heterodimer  
primarily comprised of a heavy chain of 200 kDa containing the A1, A2, and B  
25 domains and an 80 kDa light chain containing the A3, C1, and C2 domains (Kaufman  
*et al.*, J. Biol. Chem., 263:6352-6362 [1988]). Both the single chain polypeptide and  
the heterodimer circulate in the plasma as inactive precursors (Ganz *et al.*, Eur. J.  
Biochem., 170:521-528 [1988]). Activation of Factor VIII in plasma is initiated by  
thrombin cleavage between the A2 and B domains, which releases the B domain and  
30 results in a heavy chain consisting of the A1 and A2 domains. The 980 amino acid B

domain is deleted in the activated procoagulant form of the protein. Additionally, in the native protein, two polypeptide chains ("a" and "b"), flanking the B domain, are bound to a divalent calcium cation. Hemophilia may result from point mutations, deletions, or mutations resulting in a stop codon (*See, Antonarakis et al., Mol. Biol. Med.*, 4:81 [1987]).

The disease is relatively rare, afflicting approximately one in 10,000 males. Hemophilia in females is extremely rare, although it may occur in female children of an affected father and carrier mother, as well as in females with X-chromosomal abnormalities (*e.g.*, Turner syndrome, X mosaicism, etc.). The severity of each patient's disease is broadly characterized into three groups--"mild," "moderate," and "severe," depending on the severity of the patient's symptoms and circulating Factor VIII levels. While normal levels of Factor VIII range between 50 and 200 ng/mL plasma, mildly affected patients have 6-60% of this value, and moderately affected patients have 1-5% of this value. Severely affected hemophiliacs have less than 1% of normal Factor VIII levels.

While hemophiliacs clearly require clotting factor after surgery or severe trauma, on a daily basis, spontaneous internal bleeding is a greater concern. Hemophiliacs experience spontaneous hemorrhages from early infancy, as well as frequent spontaneous hemarthroses and other hemorrhages requiring clotting factor replacement. Without effective treatment, chronic hemophilic arthropathy occurs by young adulthood. Severely affected patients are prone to serious hemorrhages that may dissect through tissue planes, ultimately resulting in death due to compromised vital organs.

Hematomas are commonly observed in moderately and severely affected hemophiliacs. In these patients, hematomas have a tendency to progressively enlarge and dissect in all directions. Some of these hematomas expand locally, resulting in local compression of adjacent organs, blood vessels, and nerves. A rare, yet often fatal, complication of abdominal hematomas is the perforation and drainage of the hematoma into the colon, resulting in infection and septicemia. Intracranial and/or extracranial hemorrhage also represent very dangerous bleeding situations. While

subcutaneous hematomas may dissect into muscle, pharyngeal and retropharyngeal hematomas (*e.g.*, complicating bacterial or viral pharyngitis) may enlarge and obstruct the airway, sometimes resulting in a life-threatening situation that requires administration of a sufficient dose of Factor VIII concentrate to normalize the Factor VIII level.

In addition to hematomas, hemarthroses are commonly observed in hemophiliacs, with bleeding into the joint accounting for approximately 75% of hemophilic bleeding. Repeated hemorrhaging into the joints eventually results in extensive destruction of articular cartilage, synovial hyperplasia, and other reactive changes in adjacent tissues and bone. A major complication of repeated hemarthroses is joint deformity, which is often accompanied by muscle atrophy and soft tissue contractures; osteoporosis and cystic areas in the subchondral bone may also develop, along with progressive loss of joint space.

Other symptoms are often observed in hemophiliacs, including hematuria and mucous membrane bleeding. Hematuria is experienced by virtually all severely affected hemophiliacs sometime during their lifetimes, and mucous membrane bleeding is common in hemophiliacs. Bone cysts (pseudotumors) are rare, but dangerous complications of hemophilic bleeding. In many of these cases, immediate treatment is necessary.

In the early 1980s, many severely affected hemophiliacs were treated with Factor VIII concentrate about three times weekly. Unfortunately, these concentrates transmitted viruses, such as hepatitis B and/or C, and human immunodeficiency virus (HIV). In the United States and Western Europe, at least 75% of Factor VIII concentrate recipients have been reported to have anti-HIV antibodies (*See*, Schrier and Leung, *supra*). Some of these patients also developed HIV-associated immune thrombocytopenia, a very serious complication in hemophiliacs. In spite of antiviral therapy (*e.g.*, with zidovudine and pentamidine prophylaxis), which has tended to slow disease progression, full-blown AIDS (acquired immunodeficiency syndrome) occurs at an inexorable rate in hemophiliacs infected with HIV. Indeed, this has reversed the improvement in the life expectancy of hemophiliacs, which peaked at 66 years of age

during the 1970s, and has dropped to 49 years (*See*, Schrier and Leung, *supra*). The development of virus-free preparations and recombinant Factor VIII has helped control infectious viral contamination.

However, for hemophiliacs, the availability of viral-free concentrates and recombinant Factor VIII, while significant, is but part of the solution. In order to prevent spontaneous internal bleeding episodes, patients suffering from hemophilia A must consistently have serum Factor VIII levels of about 1%, and preferably 5%. Currently, the cost of viral-free concentrates and recombinant Factor VIII make it prohibitively expensive to administer the clotting factor prophylactically or on a maintenance basis. Indeed, most hemophiliacs in the U.S. do not receive recombinant Factor VIII therapy on a maintenance basis, but only receive it prior to activities or events which might cause bleeding (*e.g.*, surgery), or as a treatment for spontaneous bleeding.

Moreover, even if cost effective preparations of recombinant or virus-free Factor VIII were available, a steady state level of Factor VIII cannot be achieved by its daily administration. At best, patients receive widely varying levels of Factor VIII. Immediately following the administration, the levels are super-physiological, while prior to administration the levels are sub-physiological. Thus, there remains a need for methods and compositions that are relatively economic, yet effective in the treatment and prevention of bleeding in hemophiliacs, particularly spontaneous bleeds. Furthermore, there is a need in the art for methods and compositions for long term delivery of clotting factors (*e.g.*, Factor VIII) which more closely mimic the steady state physiological levels observed in normal individuals.

## SUMMARY OF THE INVENTION

The present invention provides improved viral vectors suitable for gene therapy to treat hemophilia. In particular, the present invention provides AAV vectors and methods for treating hemophilia A by delivering nucleic acids coding for the clotting protein Factor VIII. The present invention also provides pharmaceutical compositions comprising such AAV vectors, as well as methods for making and using the vectors.

The present invention is particularly suited for use in hemophilia A gene therapy. Accordingly, in one embodiment of the invention, at least one AAV vector containing a nucleic acid molecule encoding Factor VIII is operably linked to control sequences that direct expression of Factor VIII in a suitable recipient cell. The AAV  
5 vectors are then introduced into a recipient cell of the subject, under conditions that result in expression of Factor VIII. The subject, therefore, has a continuous supply of Factor VIII available to clot blood during bleeding episodes.

Using the methods of the present invention, long term expression of therapeutic levels of Factor VIII have been achieved *in vivo*. In one embodiment, animals were  
10 administered, via the portal vein, two AAV vectors: one carrying the DNA sequence coding for the heavy chain of Factor VIII and the other carrying the DNA sequence coding for the light chain of Factor VIII. Blood samples were collected periodically and assayed for Factor VIII activity. Reproducibly, animals expressed between 600 and 900 ng/ml of biologically active Factor VIII, levels that are well above the normal  
15 physiological levels of approximately 200 ng/ml. Furthermore, these levels have been sustained for over 13 months without a decrease in Factor VIII levels or activity. In a related embodiment, a B-domain deleted form of Factor VIII was cloned into a single AAV vector and shown to express biologically active Factor VIII.

It is not intended, however, that the present invention be limited to specific  
20 embodiments. Many different forms of recombinant Factor VIII have been made and tested both *in vitro* and *in vivo*, using a variety of different control and regulatory sequences. Any DNA sequence coding for biologically active Factor VIII can be expressed using the AAV vectors and methods taught in the present invention. Therefore, the present invention encompasses any AAV vector or vectors containing  
25 Factor VIII sequences that produce biologically active Factor VIII protein *in vitro* or *in vivo*.

For example, in some embodiments, the AAV vector contains the first 57 base pairs of the Factor VIII heavy chain which encodes the 10 amino acid signal sequence, as well as the human growth hormone (hGH) polyadenylation sequence. In some  
30 alternative embodiments, the vector also contains the A1 and A2 domains, as well as 5

amino acids from the N-terminus of the B domain, and/or 85 amino acids of the C-terminus of the B domain, as well as the A3, C1, and C2 domains. In yet other embodiments, the nucleic acids coding for Factor VIII heavy chain and light chain were cloned into a single vector separated by 42 nucleic acids coding for 14 amino acids of the B domain.

The present invention also provides methods for administering the above-described vectors. For example, it is intended that the present invention encompass methods suitable for delivery of the AAV vectors to the livers of recipient patients or test animals. It is not intended that the present invention be limited to any particular route of administration. However, in preferred embodiments, the AAV vectors of the present invention are successfully administered via the portal or arterial vasculature.

These and other embodiments of the invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

## **BRIEF DESCRIPTION OF THE DRAWINGS**

Figure 1 provides a schematic representation of the Factor VIII protein.

Figure 2 provides a schematic representation of a B-domain deleted form of Factor VIII protein.

Figure 3 provides a schematic representation of a B-domain deleted Factor VIII AAV construct (AAV-F8-1) from internal terminal repeat (ITR to ITR), including control sequences.

Figure 4 provides a schematic representation of a B-domain deleted Factor VIII AAV construct (PVM4.1c-F8AB) from internal terminal repeat (ITR to ITR), including control sequences.

Figure 5 provides the sequence of pAAV-F8-1 (ITR to ITR), with the plasmid backbone omitted.

Figure 6 provides the sequence of pVm4.1cF8ΔB (ITR to ITR), with the plasmid backbone omitted.

Figure 7 provides a map of rAAV-hFVIII-HC and rAAV-hFVIII-LC vectors.

Figure 8 provides a graph demonstrating the expression of various human FVIII constructs in mouse plasma.

Figure 9 provides Southern blot analyses of liver DNA using probes specific for (A) the light chain of hFVIII, and (B) the heavy chain of hFVIII.

5 Figure 10 provides Southern blot analyses of DNA from different tissues using probes specific for (A) the light chain of hFVIII, and (B) the heavy chain of hFVIII.

Figure 11 provides Northern blot analyses of liver RNA using probes specific for (A) the light chain of hFVIII, and (B) the heavy chain of hFVIII.

## 10 GENERAL DESCRIPTION OF THE INVENTION

The present invention relates to improved viral vectors useful for expressing gene products at high levels in human cells. In particular, the present invention provides AAV vectors suitable for gene therapy. These vectors are capable of delivering nucleic acid containing constructs which result in the production of Factor  
15 VIII protein in a host. The present invention also provides pharmaceutical compositions comprising such AAV vectors, as well as methods for making and using the constructs.

The AAV vectors and rAAV virions of the present invention can be produced using standard methodology known to those of skill in the art. The methods generally  
20 involve the steps of: (1) introducing an AAV vector into a host cell; (2) introducing an AAV helper construct into the host cell, where the helper construct includes AAV coding regions capable of being expressed in the host cell to complement AAV helper functions missing from the AAV vector; (3) introducing one or more helper viruses and/or accessory function vectors into the host cell, wherein the helper virus and/or  
25 accessory function vectors provide accessory functions capable of supporting efficient recombinant AAV ("rAAV") virion production in the host cell; and (4) culturing the host cell to produce rAAV virions. The AAV vector, AAV helper construct and the helper virus or accessory function vector(s) can be introduced into the host cell either simultaneously or serially, using standard transfection techniques.

Unless otherwise indicated, the practice of the present invention employs conventional methods of virology, microbiology, molecular biology and recombinant DNA techniques within the skill of the art, including those described in such references as Sambrook *et al.* (eds.) *Molecular Cloning: A Laboratory Manual*; Glover  
5 (ed.) *DNA Cloning: A Practical Approach*, Vols. I and II; Gait (ed.) *Oligonucleotide Synthesis*; Hames and Higgins (eds.) *Nucleic Acid Hybridization*; Hames and Higgins (eds.) *Transcription and Translation*; Tijessen (ed.) *CRC Handbook of Parvoviruses*, Vols. I and II; and Fields and Knipe (eds.) *Fundamental Virology*, 2nd Edition, Vols. I and II.

### Definitions

In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below.

As used herein, the terms "gene transfer" and "gene delivery" refer to methods  
15 or systems for reliably inserting a particular nucleotide sequence (*e.g.*, DNA) into targeted cells. In particularly preferred embodiments, the nucleotide sequence comprises at least a portion of Factor VIII.

As used herein, the terms "vector," and "gene transfer vector" refer to any genetic element, such as a plasmid, phage, transposon, cosmid, chromosome, virus,  
20 virion, etc., which is capable of replication when associated with the proper control sequences and/or which can transfer nucleic acid sequences between cells. Thus, the term includes cloning and expression vectors, as well as viral vectors.

Gene transfer vectors may include transcription sequences such as polyadenylation sites, selectable markers or reporter genes, enhancer sequences, and  
25 other control sequences which allow for the induction of transcription. Such control sequences are described more fully below.

The term "expression vector" as used herein refers to a recombinant DNA molecule containing a desired coding sequence and appropriate nucleic acid sequences necessary for the expression of the operably linked coding sequence in a particular

host organism. Nucleic acid sequences necessary for expression in prokaryotes usually include a promoter, an operator (optional), and a ribosome binding site, as well as other sequences. Eukaryotic cells are generally known to utilize promoters (constitutive, inducible or tissue specific), enhancers, and termination and polyadenylation signals, although some elements may be deleted and other elements added without sacrificing the necessary expression.

As used herein, the terms "host" and "expression host" refer to organisms and/or cells which harbor an exogenous DNA sequence (*e.g.*, via transfection), an expression vector or vehicle, as well as organisms and/or cells that are suitable for use in expressing a recombinant gene or protein. It is not intended that the present invention be limited to any particular type of cell or organism. Indeed, it is contemplated that any suitable organism and/or cell will find use in the present invention as a host.

As used herein, the terms "viral replicons" and "viral origins of replication" refer to viral DNA sequences that allow for the extrachromosomal replication of a vector in a host cell expressing the appropriate replication factors. In some embodiments, vectors which contain either the SV40 or polyoma virus origin of replication replicate to high copy number, while vectors which contain the replicons from bovine papillomavirus or Epstein-Barr virus replicate extrachromosomally at low copy number may be utilized in other embodiments.

As used herein, the term "AAV vector" refers to a vector having functional or partly functional ITR sequences. The ITR sequences may be derived from an adeno-associated virus serotype, including without limitation, AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-X7, etc. The ITRs, however, need not be the wild-type nucleotide sequences, and may be altered (*e.g.*, by the insertion, deletion or substitution of nucleotides), so long as the sequences retain function provide for functional rescue, replication and packaging. AAV vectors can have one or more of the AAV wild-type genes deleted in whole or part, preferably the *rep* and/or *cap* genes but retain functional flanking ITR sequences. Functional ITR sequences are necessary for the rescue, replication and packaging of the AAV virion. Thus, an "AAV vector" is

defined herein to include at least those sequences required in *cis* for replication and packaging (e.g., functional ITRs) of the virus.

As used herein, the term "ITR" refers to inverted terminal repeats. The terms "adeno-associated virus inverted terminal repeats" or "AAV ITRs" refer to the art-recognized palindromic regions found at each end of the AAV genome which function together in *cis* as origins of DNA replication and as packaging signals for the virus. For use in some embodiments of the present invention, flanking AAV ITRs are positioned 5' and 3' of one or more selected heterologous nucleotide sequences. Optionally, the ITRs together with the *rep* coding region or the Rep expression product provide for the integration of the selected sequences into the genome of a target cell.

As used herein, the term "AAV *rep* coding region" refers to the art-recognized region of the AAV genome which encodes the replication proteins Rep 78, Rep 68, Rep 52 and Rep 40. These Rep expression products have been shown to possess many functions, including recognition, binding and nicking of the AAV origin of DNA replication, DNA helicase activity and modulation of transcription from AAV (or other heterologous) promoters. The Rep expression products are collectively required for replicating the AAV genome. Muzyczka (Muzyczka, Curr. Top. Microbiol. Immunol., 158:97-129 [1992]) and Kotin (Kotin, Hum. Gene Ther., 5:793-801 [1994]) provide additional descriptions of the AAV *rep* coding region, as well as the *cap* coding region described below. Suitable homologues of the AAV *rep* coding region include the human herpesvirus 6 (HHV-6) *rep* gene which is also known to mediate AAV-2 DNA replication (Thomson *et al.*, Virol., 204:304-311 [1994]).

As used herein, the term "AAV *cap* coding region" refers to the art-recognized region of the AAV genome which encodes the capsid proteins VP1, VP2, and VP3, or functional homologues thereof. These *cap* expression products supply the packaging functions which are collectively required for packaging the viral genome.

As used herein, the term "AAV helper function" refers to AAV coding regions capable of being expressed in the host cell to complement AAV viral functions missing from the AAV vector. Typically, the AAV helper functions include the AAV

*rep* coding region and the AAV *cap* coding region. An "AAV helper construct" is a vector containing AAV coding regions required to complement AAV viral functions missing from the AAV vector (*e.g.*, the AAV *rep* coding region and the AAV *cap* coding region).

5 As used herein, the terms "accessory functions" and "accessory factors" refer to functions and factors that are required by AAV for replication, but are not provided by the AAV virion (or rAAV virion) itself. Thus, these accessory functions and factors must be provided by the host cell, a virus (*e.g.*, adenovirus or herpes simplex virus), or another expression vector that is co-expressed in the same cell. Generally, the E1,  
10 E2A, E4 and VA coding regions of adenovirus are used to supply the necessary accessory function required for AAV replication and packaging (Matsushita *et al.*, Gene Therapy 5:938 [1998]).

As used herein, the term "wild type" ("wt") refers to a gene or gene product which has the characteristics of that gene or gene product when isolated from a  
15 naturally occurring source. A wild-type gene is that which is most frequently observed in a population and is thus arbitrarily designed the "normal" or "wild-type" form of the gene. In contrast, the term "modified" or "mutant" refers to a gene or gene product which displays modifications in sequence and or functional properties (*i.e.*, altered characteristics) when compared to the wild-type gene or gene product. It is noted that  
20 naturally-occurring mutants can be isolated; these are identified by the fact that they have altered characteristics when compared to the wild-type gene or gene product.

As used herein, the term "AAV virion" refers to a complete virus particle, such as a "wild-type" (wt) AAV virus particle (comprising a linear, single-stranded AAV nucleic acid genome associated with an AAV capsid protein coat). In this regard,  
25 single-stranded AAV nucleic acid molecules of either complementary sense (*e.g.*, "sense" or "antisense" strands), can be packaged into any one AAV virion and both strands are equally infectious.

As used herein, the terms "recombinant AAV virion," and "rAAV virion" refer to an infectious viral particle containing a heterologous DNA molecule of interest (*e.g.*,

Factor VIII sequence) which is flanked on both sides by AAV ITRs. In some embodiments of the present invention, an rAAV virion is produced in a suitable host cell which contains an AAV vector, AAV helper functions and accessory functions introduced therein. In this manner, the host cell is rendered capable of encoding AAV polypeptides that are required for packaging the AAV vector containing a recombinant nucleotide sequence of interest, such as at least a portion of Factor VIII or portions of Factor VIII domains, into recombinant virion particles for subsequent gene delivery.

As used herein, the term "transfection" refers to the uptake of foreign DNA by a cell, and a cell has been "transfected" when exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are generally known in the art (See e.g., Graham *et al.*, *Virol.*, 52:456 [1973]; Sambrook *et al.*, *Molecular Cloning, a Laboratory Manual*, Cold Spring Harbor Laboratories, New York [1989]; Davis *et al.*, *Basic Methods in Molecular Biology*, Elsevier, [1986]; and Chu *et al.*, *Gene* 13:197 [1981]). Such techniques can be used to introduce one or more exogenous DNA moieties, such as a gene transfer vector and other nucleic acid molecules, into suitable recipient cells.

As used herein, the terms "stable transfection" and "stably transfected" refers to the introduction and integration of foreign DNA into the genome of the transfected cell. The term "stable transfectant" refers to a cell which has stably integrated foreign DNA into the genomic DNA.

As used herein, the term "transient transfection" or "transiently transfected" refers to the introduction of foreign DNA into a cell where the foreign DNA fails to integrate into the genome of the transfected cell. The foreign DNA persists in the nucleus of the transfected cell for several days. During this time the foreign DNA is subject to the regulatory controls that govern the expression of endogenous genes in the chromosomes. The term "transient transfectant" refers to cells which have taken up foreign DNA but have failed to integrate this DNA.

As used herein, the term "transduction" denotes the delivery of a DNA molecule to a recipient cell either *in vivo* or *in vitro*, via a replication-defective viral vector, such as via a recombinant AAV virion.

As used herein, the term "recipient cell" refers to a cell which has been transfected or transduced, or is capable of being transfected or transduced, by a nucleic acid construct or vector bearing a selected nucleotide sequence of interest (*i.e.*, Factor VIII). The term includes the progeny of the parent cell, whether or not the progeny are identical in morphology or in genetic make-up to the original parent, so long as the selected nucleotide sequence is present.

The term "heterologous" as it relates to nucleic acid sequences such as coding sequences and control sequences, denotes sequences that are not normally joined together, and/or are not normally associated with a particular cell. Thus, a "heterologous" region of a nucleic acid construct or a vector is a segment of nucleic acid within or attached to another nucleic acid molecule that is not found in association with the other molecule in nature. For example, a heterologous region of a nucleic acid construct could include a coding sequence flanked by sequences not found in association with the coding sequence in nature. Another example of a heterologous coding sequence is a construct where the coding sequence itself is not found in nature (*e.g.*, synthetic sequences having codons different from the native gene). Similarly, a cell transfected with a construct which is not normally present in the cell would be considered heterologous for purposes of this invention. Allelic variation or naturally occurring mutational events do not give rise to heterologous DNA, as used herein.

As used herein, "coding sequence" or a sequence which "encodes" a particular antigen, is a nucleic acid sequence which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide *in vitro* or *in vivo*, when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A coding sequence can include, but is not limited to, cDNA from prokaryotic or eukaryotic mRNA, genomic DNA sequences from prokaryotic or eukaryotic DNA, and even synthetic DNA sequences. A transcription termination sequence will usually be located 3' to the coding sequence.

As used herein, the term "nucleic acid" sequence refers to a DNA or RNA sequence. The term captures sequences that include any of the known base analogues

of DNA and RNA such as, but not limited to 4-acetylcytosine, 8-hydroxy-N6-methyladenosine, aziridinylcytosine, pseudoisocytosine, 5-(carboxyhydroxymethyl)uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethylaminomethyluracil, dihydrouracil, inosine, N6-isopentenyladenine, 1-methyladenine, 1-methylpseudouracil, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarbonylmethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, N-uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine.

As used herein, the term "recombinant DNA molecule" as used herein refers to a DNA molecule which is comprised of segments of DNA joined together by means of molecular biological techniques.

As used herein, the term "regulatory element" refers to a genetic element which controls some aspect of the expression of nucleic acid sequences. For example, a promoter is a regulatory element which facilitates the initiation of transcription of an operably linked coding region. Other regulatory elements are splicing signals, polyadenylation signals, termination signals, etc. (defined *infra*).

The term DNA "control sequences" refers collectively to regulatory elements such as promoter sequences, polyadenylation signals, transcription termination sequences, upstream regulatory domains, origins of replication, internal ribosome entry sites ("IRES"), enhancers, and the like, which collectively provide for the replication, transcription and translation of a coding sequence in a recipient cell. Not all of these control sequences need always be present so long as the selected coding sequence is capable of being replicated, transcribed and translated in an appropriate recipient cell.

Transcriptional control signals in eukaryotes generally comprise "promoter" and "enhancer" elements. Promoters and enhancers consist of short arrays of DNA sequences that interact specifically with cellular proteins involved in transcription (Maniatis *et al.*, Science 236:1237 [1987]). Promoter and enhancer elements have  
5 been isolated from a variety of eukaryotic sources including genes in yeast, insect and mammalian cells and viruses (analogous control sequences, *i.e.*, promoters, are also found in prokaryotes). The selection of a particular promoter and enhancer depends on what cell type is to be used to express the protein of interest (*i.e.*, Factor VIII). Some eukaryotic promoters and enhancers have a broad host range while others are  
10 functional in a limited subset of cell types (*See e.g.*, Voss *et al.*, Trends Biochem. Sci., 11:287 [1986]; and Maniatis *et al.*, *supra*, for reviews). For example, the SV40 early gene enhancer is very active in a wide variety of cell types from many mammalian species and has been widely used for the expression of proteins in mammalian cells (Dijkema *et al.*, EMBO J. 4:761 [1985]). Two other examples of promoter and  
15 enhancer elements active in a broad range of mammalian cell types are those from the human elongation factor 1 $\alpha$  gene (Uetsuki *et al.*, J. Biol. Chem., 264:5791 [1989]; Kim *et al.*, Gene 91:217 [1990]; and Mizushima and Nagata, Nucl. Acids. Res., 18:5322 [1990]) and the long terminal repeats of the Rous sarcoma virus (Gorman *et al.*, Proc. Natl. Acad. Sci. USA 79:6777 [1982]) and the human cytomegalovirus  
20 (Boshart *et al.*, Cell 41:521 [1985]). Promoters and enhancers can be found naturally alone or together. For example, the long terminal repeats of retroviruses contain both promoter and enhancer functions. Moreover, generally promoters and enhancers act independently of the gene being transcribed or translated. Thus, the enhancer and promoter may be "endogenous" or "exogenous" or "heterologous." An "endogenous"  
25 enhancer/promoter is one which is naturally linked with a given gene in the genome. An "exogenous" or "heterologous" enhancer and promoter is one which is placed in juxtaposition to a gene by means of genetic manipulation (*i.e.*, molecular biological techniques) such that transcription of that gene is directed by the linked enhancer/promoter.

As used herein, the term "tissue specific" refers to regulatory elements or control sequences, such as a promoter, enhancers, etc., wherein the expression of the nucleic acid sequence is substantially greater in a specific cell type(s) or tissue(s). In particularly preferred embodiments, the albumin promoter and the transthyretin promoter display increased expression of FVIII in hepatocytes, as compared to other cell types. It is not intended, however, that the present invention be limited to the albumin or transthyretin promoters or to hepatic-specific expression, as other tissue specific regulatory elements, or regulatory elements that display altered gene expression patterns, are contemplated.

The presence of "splicing signals" on an expression vector often results in higher levels of expression of the recombinant transcript. Splicing signals mediate the removal of introns from the primary RNA transcript and consist of a splice donor and acceptor site (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, New York [1989], pp. 16.7-16.8). A commonly used splice donor and acceptor site is the splice junction from the 16S RNA of SV40.

Efficient expression of recombinant DNA sequences in eukaryotic cells requires expression of signals directing the efficient termination and polyadenylation of the resulting transcript. Transcription termination signals are generally found downstream of the polyadenylation signal and are a few hundred nucleotides in length. The term "poly A site" or "poly A sequence" as used herein denotes a DNA sequence which directs both the termination and polyadenylation of the nascent RNA transcript. Efficient polyadenylation of the recombinant transcript is desirable as transcripts lacking a poly A tail are unstable and are rapidly degraded. The poly A signal utilized in an expression vector may be "heterologous" or "endogenous." An endogenous poly A signal is one that is found naturally at the 3' end of the coding region of a given gene in the genome. A heterologous poly A signal is one which is isolated from one gene and placed 3' of another gene. A commonly used heterologous poly A signal is the SV40 poly A signal. The SV40 poly A signal is contained on a 237 bp *Bam*HI/*Bcl*II restriction fragment and directs both termination and polyadenylation (Sambrook *et al.*, *supra*, at 16.6-16.7).

"Operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, control sequences operably linked to a coding sequence are capable of effecting the expression of the coding sequence. The control sequences need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

The term "isolated" when used in relation to a nucleic acid, as in "an isolated oligonucleotide" or "isolated polynucleotide" refers to a nucleic acid sequence that is identified and separated from at least one contaminant nucleic acid with which it is ordinarily associated in its natural source. Isolated nucleic acid is such present in a form or setting that is different from that in which it is found in nature. In contrast, non-isolated nucleic acids are nucleic acids such as DNA and RNA found in the state they exist in nature. For example, a given DNA sequence (*e.g.*, a gene) is found on the host cell chromosome in proximity to neighboring genes; RNA sequences, such as a specific mRNA sequence encoding a specific protein, are found in the cell as a mixture with numerous other mRNAs which encode a multitude of proteins. The isolated nucleic acid, oligonucleotide, or polynucleotide may be present in single-stranded or double-stranded form. When an isolated nucleic acid, oligonucleotide or polynucleotide is to be utilized to express a protein, the oligonucleotide or polynucleotide will contain at a minimum the sense or coding strand (*i.e.*, the oligonucleotide or polynucleotide may single-stranded), but may contain both the sense and anti-sense strands (*i.e.*, the oligonucleotide or polynucleotide may be double-stranded).

As used herein, the term "purified" or "to purify" refers to the removal of contaminants from a sample. For example, antibodies may be purified by removal of contaminating non-immunoglobulin proteins; they may also purified by the removal of immunoglobulin that does not bind the antigen of interest (*e.g.*, at least a portion of Factor VIII). The removal of non-immunoglobulin proteins and/or the removal of

immunoglobulins that do not bind the antigen of interest (*e.g.*, at least a portion of Factor VIII) results in an increase in the percent of desired antigen-reactive immunoglobulins in the sample. In another example, recombinant polypeptides of Factor VIII are expressed in bacterial host cells and the polypeptides are purified by the removal of host cell proteins; the percent of recombinant polypeptides is thereby increased in the sample.

As used herein, the term "chimeric protein" refers to two or more coding sequences obtained from different genes, that have been cloned together and that, after translation, act as a single polypeptide sequence. Chimeric proteins are also referred to as "hybrid proteins." As used herein, the term "chimeric protein" refers to coding sequences that are obtained from different species of organisms, as well as coding sequences that are obtained from the same species of organisms.

A "composition comprising a given polynucleotide sequence" as used herein refers broadly to any composition containing the given polynucleotide sequence. The composition may comprise an aqueous solution.

As used herein, the term "at risk" is used in references to individuals who are at risk for experiencing hemorrhagic episodes. In particularly preferred embodiments, the individuals are hemophiliacs with mild, moderate, or severe hemophilia.

As used herein, the term "subject" refers to any animal (*i.e.*, vertebrates and invertebrates), while the term "vertebrate subject" refers to any member of the subphylum Chordata. It is intended that the term encompass any member of this subphylum, including, but not limited to humans and other primates, rodents (*e.g.*, mice, rats, and guinea pigs), lagamorphs (*e.g.*, rabbits), bovines (*e.g.*, cattle), ovines (*e.g.*, sheep), caprines (*e.g.*, goats), porcines (*e.g.*, swine), equines (*e.g.*, horses), canines (*e.g.*, dogs), felines (*e.g.*, cats), domestic fowl (*e.g.*, chickens, turkeys, ducks, geese, other gallinaceous birds, etc.), as well as feral or wild animals, including, but not limited to, such animals as ungulates (*e.g.*, deer), bear, fish, lagamorphs, rodents, birds, etc. It is not intended that the term be limited to a particular age or sex. Thus,

adult and newborn subjects, as well as fetuses, whether male or female, are encompassed by the term.

As defined herein, a "therapeutically effective amount" or "therapeutic effective dose" is an amount or dose of AAV vector or virions capable of producing sufficient amounts of Factor VIII to decrease the time it takes for a subject's blood to clot. Generally, severe hemophiliacs having less than 1% of normal levels of FVIII have a whole blood clotting time of greater than 60 minutes as compared to approximately 10 minutes for non-hemophiliacs.

## 10 DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to AAV vectors suitable for hemophilia A gene therapy. More particularly, these AAV vectors are suitable for delivering nucleic acids encoding Factor VIII into a recipient host suspected of suffering from a blood clotting disorder. Using the nucleic acid as a template, the host produces Factor VIII, such that the subject's blood is able to clot. The present invention also provides pharmaceutical compositions comprising such AAV vectors, as well as methods for making and using the constructs.

### I. AAV Vectors

Adeno-associated virus (AAV) is a non-pathogenic, replication-defective, helper-dependent parvovirus (or "dependovirus" or "adeno-satellite virus"). There are at least six recognized serotypes, designated as AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-X7, etc. Culture and serologic evidence indicates that human infection occurs with AAV-2 and AAV-3. Although 85% of the human population is seropositive for AAV-2, the virus has never been associated with disease in humans. Recombinant AAV (rAAV) virions are of interest as vectors for gene therapy because of their broad host range, excellent safety profile, and duration of transgene expression in infected hosts. One remarkable feature of recombinant AAV (rAAV) virions is the prolonged expression achieved after *in vivo* administration.

AAV vectors of the present invention may be constructed using known techniques to provide, as operatively linked components in the direction of transcription, (a) control sequences including a transcriptional initiation and termination regions, and (b) a nucleotide sequence encoding at least a portion of Factor VIII. The control sequences are selected to be functional in a targeted recipient cell. The resulting construct which contains the operatively linked components is bounded (5' and 3') with functional AAV ITR sequences.

The nucleotide sequences of AAV ITR regions are known (*See e.g.*, Kotin, Hum. Gene Ther., 5:793-801 [1994]; Berns, "Parvoviridae and Their Replication" in Fields and Knipe (eds), *Fundamental Virology*, 2nd Edition, for the AAV-2 sequence). AAV ITRs used in the vectors of the invention need not have a wild-type nucleotide sequence, and may be altered (*e.g.*, by the insertion, deletion or substitution of nucleotides). Additionally, AAV ITRs may be derived from any of several AAV serotypes, including without limitation, AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAVX7, etc. Furthermore, 5' and 3' ITRs which flank a selected nucleotide sequence in an AAV vector need not necessarily be identical or derived from the same AAV serotype or isolate, so long as they function as intended.

#### A. Control Sequences

In some embodiments of the present invention, heterologous control sequences are employed with the vectors. Useful heterologous control sequences generally include those derived from sequences encoding mammalian or viral genes. Examples include, but are not limited to, the SV40 early promoter, mouse mammary tumor virus LTR promoter, adenovirus major late promoter (Ad MLP), a herpes simplex virus (HSV) promoter, a cytomegalovirus (CMV) promoter such as the CMV immediate early promoter region (CMVIE), a rous sarcoma virus (RSV) promoter, synthetic promoters, hybrid promoters, and the like. In addition, sequences derived from nonviral genes, such as the murine metallothionein gene, also find use herein. Such promoter sequences are commercially available (*e.g.*, from Stratagene).

It is contemplated that in some embodiments, tissue-specific expression may be desirable (*e.g.*, expression of biologically active Factor VIII by hepatocytes). It is not intended that the present invention be limited to expression of biologically active Factor VIII by any particular cells or cell types. However, as hepatocytes (*i.e.*, liver cells) are the cells that normally synthesized Factor VIII (*See*, Kaufman, *Ann. Rev. Med.*, 43:325 [1992]), it is contemplated that in some particularly preferred embodiments, the compositions of the present invention be administered to the liver.

In preferred embodiments, expression is achieved by coupling the coding sequence for Factor VIII with heterologous control sequences derived from genes that are specifically transcribed by a selected tissue type. A number of tissue-specific promoters have been described above which enable directed expression in selected tissue types. However, control sequences used in the present AAV vectors can also comprise control sequences normally associated with the selected nucleic acid sequences.

#### **B. Construction of AAV Factor VIII Vectors**

AAV vectors that contain a control sequence and a nucleotide sequence of interest (*i.e.*, at least a portion of the sequence encoding Factor VIII), bounded by AAV ITRs (*i.e.*, AAV vectors), can be constructed by directly inserting selected sequences into an AAV genome with the major AAV open reading frames ("ORFs") excised. Other portions of the AAV genome can also be deleted, so long as a sufficient portion of the ITRs remain to allow for replication and packaging functions. These constructs can be designed using techniques well known in the art (*See e.g.*, U.S. Patent Nos. 5,173,414 and 5,139,941, all of which are herein incorporated by reference); International Publication Nos. WO 92/01070 and WO 93/03769; Lebkowski *et al.*, *Mol. Cell. Biol.*, 8:3988-3996 [1988]; Vincent *et al.*, *Vaccines 90* [Cold Spring Harbor Laboratory Press, 1990]; Carter, *Curr. Opin. Biotechnol.*, 3:533-539 [1992]; Muzyczka, *Curr. Top. Microbiol. Immunol.*, 158:97-129 [1992]; Kotin, *Hum. Gene*

Ther., 5:793-801 [1994]; Shelling and Smith, Gene Ther., 1:165-169 [1994]; and Zhou *et al.*, J. Exp. Med., 179:1867-1875 [1994]).

Alternatively, AAV ITRs can be excised from the viral genome or from an AAV vector containing the same and fused 5' and 3' of a selected nucleic acid construct that is present in another vector using standard ligation techniques, such as those described in Sambrook *et al.*, *supra*. For example, ligations can be accomplished in 20 mM Tris-Cl pH 7.5, 10 mM MgCl<sub>2</sub>, 10 mM DTT, 33 µg/ml BSA, 10 mM-50 mM NaCl, and either 40 µM ATP, 0.01-0.02 (Weiss) units T4 DNA ligase at 0°C (for "sticky end" ligation) or 1 mM ATP, 0.3-0.6 (Weiss) units T4 DNA ligase at 14°C (for "blunt end" ligation). Intermolecular "sticky end" ligations are usually performed at 30-100 µg/ml total DNA concentrations (5-100 nM total end concentration). AAV vectors which contain ITRs have been described in (*e.g.*, U.S. Patent No. 5,139,941, herein incorporated by reference). In particular, several AAV vectors are described therein which are available from the American Type Culture Collection ("ATCC") under Accession Numbers 53222, 53223, 53224, 53225 and 53226.

Additionally, chimeric genes can be produced synthetically to include AAV ITR sequences arranged 5' and 3' of a selected nucleic acid sequence. The complete chimeric sequence is assembled from overlapping oligonucleotides prepared by standard methods (*See e.g.*, Edge, Nature 292:756 [1981]; Nambair *et al.*, Science 223:1299 [1984]; and Jay *et al.*, J. Biol. Chem., 259:6311 [1984]).

Moreover, it is not intended that the present invention be limited to any specific Factor VIII sequence. Many natural and recombinant forms of Factor VIII have been isolated and assayed both *in vitro* and *in vivo*, using a variety of different regulatory elements and control sequences. Therefore, any known, or later discovered, DNA sequence coding for biologically active Factor VIII can be expressed, alone or in combination with at least one additional vector, using the AAV vectors and methods taught in the present invention. Examples of naturally occurring and recombinant forms of Factor VIII can be found in the patent and scientific literature including, U.S. 5,563,045, U.S. 5,451,521, U.S. 5,422,260, U.S. 5,004,803, U.S. 4,757,006, U.S.

5,661,008, U.S. 5,789,203, U.S. 5,681,746, U.S. 5,595,886, U.S. 5,045,455, U.S.  
5,668,108, U.S. 5,633,150, U.S. 5,693,499, U.S. 5,587,310, U.S. 5,171,844, U.S.  
5,149,637, U.S. 5,112,950, U.S. 4,886,876, WO 94/11503, WO 87/07144, WO  
92/16557, WO 91/09122, WO 97/03195, WO 96/21035, WO 91/07490, EP 0 672  
5 138, EP 0 270 618, EP 0 182 448, EP 0 162 067, EP 0 786 474, EP 0 533 862,  
EP 0 506 757, EP 0 874 057, EP 0 795 021, EP 0 670 332, EP 0 500 734, EP 0  
232 112, EP 0 160 457, Sanberg *et al.*, XXth Int. Congress of the World Fed. Of  
Hemophilia (1992), and Lind *et al.*, Eur. J. Biochem., 232:19 (1995).

Nucleic acid sequences coding for the above-described Factor VIII can be  
10 obtained using recombinant methods, such as by screening cDNA and genomic  
libraries from cells expressing Factor VIII or by deriving the sequence from a vector  
known to include the same. Furthermore, the desired sequence can be isolated directly  
from cells and tissues containing the same, using standard techniques, such as phenol  
extraction and PCR of cDNA or genomic DNA (*See e.g.*, Sambrook *et al.*, *supra*, for a  
15 description of techniques used to obtain and isolate DNA). Nucleotide sequences  
encoding an antigen of interest (*i.e.*, Factor VIII sequence) can also be produced  
synthetically, rather than cloned. The complete sequence can be assembled from  
overlapping oligonucleotides prepared by standard methods and assembled into a  
complete coding sequence (*See e.g.*, Edge, Nature 292:756 [1981]; Nambair *et al.*, Sci-  
20 ence 223:1299 [1984]; and Jay *et al.*, J. Biol. Chem., 259:6311 [1984]).

Although it is not intended that the present invention be limited to any  
particular methods for assessing the production of biologically active Factor VIII, such  
methods as immunoassays (*e.g.*, ELISA) and biological activity assays are  
contemplated (*e.g.*, coagulation activity assays).

25 Furthermore, while in particularly preferred embodiments, human Factor VIII is  
encompassed by the present invention, it is not intended that the present invention be  
limited to human Factor VIII. Indeed, it is intended that the present invention  
encompass Factor VIII from animals other than humans, including but not limited to  
companion animals (*e.g.*, canines, felines, and equines), livestock (*e.g.*, bovines,

caprines, and ovines), laboratory animals (e.g., rodents such as murines, as well as lagamorphs), and "exotic" animals (e.g., marine mammals, large cats, etc.).

## II. Virion Production

5 Producing AAV Factor VIII vectors and rAAV Factor VIII virions of the present invention generally involve the steps of: (1) introducing an AAV vector containing the Factor VIII gene into a host cell; (2) introducing an AAV helper construct into the host cell, where the helper construct includes AAV coding regions capable of being expressed in the host cell to complement AAV helper functions  
10 missing from the AAV vector; (3) introducing one or more helper viruses and/or accessory function vectors into the host cell, wherein the helper virus and/or accessory function vectors provide accessory functions capable of supporting efficient recombinant AAV ("rAAV") virion production in the host cell; and (4) culturing the host cell to produce rAAV virions.

15 The above-described vectors and constructs can be introduced into a cell using standard methodology known to those of skill in the art (e.g., transfection). A number of transfection techniques are generally known in the art (*See e.g., Graham et al., Virol., 52:456 [1973], Sambrook et al. supra, Davis et al., supra, and Chu et al., Gene 13:197 [1981]*). Particularly suitable transfection methods include calcium phosphate  
20 co-precipitation (*Graham et al., Virol., 52:456-467 [1973]*), direct micro-injection into cultured cells (*Capecchi, Cell 22:479-488 [1980]*), electroporation (*Shigekawa et al., BioTechn., 6:742-751 [1988]*), liposome-mediated gene transfer (*Mannino et al., BioTechn., 6:682-690 [1988]*), lipid-mediated transduction (*Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 [1987]*), and nucleic acid delivery using high-velocity  
25 microprojectiles (*Klein et al., Nature 327:70-73 [1987]*).

For the purposes of the invention, suitable host cells for producing rAAV virions include microorganisms, yeast cells, insect cells, and mammalian cells, that can be, or have been, used as recipients of a heterologous DNA molecule. The term includes the progeny of the original cell which has been transfected. Thus, as

indicated above, a "host cell" as used herein generally refers to a cell which has been transfected with an exogenous DNA sequence. Cells from the stable human cell line, 293 (ATCC Accession No. CRL1573) are preferred in the practice of the present invention. Particularly, the human cell line 293 is a human embryonic kidney cell line  
5 that has been transformed with adenovirus type-5 DNA fragments (Graham *et al.*, J. Gen. Virol., 36:59 [1977]), and expresses the adenoviral E1a and E1b genes (Aiello *et al.*, Virol., 94:460 [1979]). The 293 cell line is readily transfected, and provides a particularly convenient platform in which to produce rAAV virions.

Host cells containing the above-described AAV vectors must be rendered  
10 capable of providing AAV helper functions in order to replicate and encapsidate the nucleotide sequences flanked by the AAV ITRs to produce rAAV virions. AAV helper functions are generally AAV-derived coding sequences which can be expressed to provide AAV gene products that, in turn, function in *trans* for productive AAV replication. AAV helper functions are used herein to complement necessary AAV  
15 functions that are missing from the AAV vectors. Thus, AAV helper functions include one, or both of the major AAV ORFs, namely the *rep* and *cap* coding regions, or functional homologues thereof.

AAV helper functions are introduced into the host cell by transfecting the host cell with an AAV helper construct either prior to, or concurrently with, the transfection  
20 of the AAV vector. AAV helper constructs are thus used to provide at least transient expression of AAV *rep* and/or *cap* genes to complement missing AAV functions that are necessary for productive AAV infection. AAV helper constructs lack AAV ITRs and can neither replicate nor package themselves.

In preferred embodiments, these constructs are in the form of a vector,  
25 including, but not limited to, plasmids, phages, transposons, cosmids, viruses, or virions. A number of AAV helper constructs have been described, such as the commonly used plasmids pAAV/Ad and pIM29+45 which encode both Rep and Cap expression products (See *e.g.*, Samulski *et al.*, J. Virol., 63:3822-3828 [1989]; and McCarty *et al.*, J. Virol., 65:2936-2945 [1991]). A number of other vectors have been

described which encode Rep and/or Cap expression products (*See e.g.*, U.S. Patent No. 5,139,941, herein incorporated by reference).

Both AAV vectors and AAV helper constructs can be constructed to contain one or more optional selectable markers. Suitable markers include genes which confer antibiotic resistance or sensitivity to, impart color to, or change the antigenic characteristics of those cells which have been transfected with a nucleic acid construct containing the selectable marker when the cells are grown in an appropriate selective medium. Several selectable marker genes that are useful in the practice of the invention include the gene encoding aminoglycoside phosphotransferase (APH) that allows selection in mammalian cells by conferring resistance to G418 (Sigma). Other suitable markers are known to those of skill in the art.

The host cell (or packaging cell) must also be rendered capable of providing non-AAV derived functions, or "accessory functions," in order to produce rAAV virions. Accessory functions are non-AAV derived viral and/or cellular functions upon which AAV is dependent for its replication. Thus, accessory functions include at least those non-AAV proteins and RNAs that are required in AAV replication, including those involved in activation of AAV gene transcription, stage specific AAV mRNA splicing, AAV DNA replication, synthesis of *rep* and *cap* expression products and AAV capsid assembly. Viral-based accessory functions can be derived from any of the known helper viruses.

Particularly, accessory functions can be introduced into and then expressed in host cells using methods known to those of skill in the art. Commonly, accessory functions are provided by infection of the host cells with an unrelated helper virus. A number of suitable helper viruses are known, including adenoviruses; herpesviruses such as herpes simplex virus types 1 and 2; and vaccinia viruses. Nonviral accessory functions will also find use herein, such as those provided by cell synchronization using any of various known agents (*See e.g.*, Buller *et al.*, J. Virol., 40:241-247 [1981]; McPherson *et al.*, Virol., 147:217-222 [1985]; and Schlehofer *et al.*, Virol., 152:110-117 [1986]).

Alternatively, accessory functions can be provided using an accessory function vector. Accessory function vectors include nucleotide sequences that provide one or more accessory functions. An accessory function vector is capable of being introduced into a suitable host cell in order to support efficient AAV virion production in the host cell. Accessory function vectors can be in the form of a plasmid, phage, virus, transposon or cosmid. Accessory vectors can also be in the form of one or more linearized DNA or RNA fragments which, when associated with the appropriate control sequences and enzymes, can be transcribed or expressed in a host cell to provide accessory functions.

Nucleic acid sequences providing the accessory functions can be obtained from natural sources, such as from the genome of adenovirus, or constructed using recombinant or synthetic methods known in the art. In this regard, adenovirus-derived accessory functions have been widely studied, and a number of adenovirus genes involved in accessory functions have been identified and partially characterized (*See e.g.*, Carter, "Adeno-Associated Virus Helper Functions," in *CRC Handbook of Parvoviruses*, Vol. I (P. Tijssen, ed.) [1990], and Muzyczka, *Curr. Top. Microbiol. Immun.*, 158:97-129 [1992]). Specifically, early adenoviral gene regions E1a, E2a, E4, VAI RNA and, possibly, E1b are thought to participate in the accessory process (Janik *et al.*, *Proc. Natl. Acad. Sci. USA* 78:1925-1929 [1981]). Herpesvirus-derived accessory functions have been described (*See e.g.*, Young *et al.*, *Prog. Med. Virol.*, 25:113 [1979]). Vaccinia virus-derived accessory functions have also been described (*See e.g.*, Carter, *supra.*, and Schlehofer *et al.*, *Virol.*, 152:110-117 [1986]).

As a consequence of the infection of the host cell with a helper virus, or transfection of the host cell with an accessory function vector, accessory functions are expressed which transactivate the AAV helper construct to produce AAV Rep and/or Cap proteins. The Rep expression products direct excision of the recombinant DNA (including the DNA of interest encoding at least a portion of Factor VIII) from the AAV vector. The Rep proteins also serve to duplicate the AAV genome. The expressed Cap proteins assemble into capsids, and the recombinant AAV genome is

packaged into the capsids. Thus, productive AAV replication ensues, and the DNA is packaged into rAAV virions.

Following recombinant AAV replication, rAAV virions can be purified from the host cell using a variety of conventional purification methods, such as CsCl  
5 gradients. Further, if helper virus infection is employed to express the accessory functions, residual helper virus can be inactivated, using known methods. For example, adenovirus can be inactivated by heating to temperatures of approximately 60°C for approximately 20 minutes or more, as appropriate. This treatment selectively inactivates the helper adenovirus which is heat labile, while preserving the rAAV  
10 which is heat stable.

### III. Pharmaceutical Compositions

The resulting rAAV virions are then ready for use in pharmaceutical compositions which can be delivered to a subject, so as to allow production of  
15 biologically active Factor VIII. Pharmaceutical compositions comprise sufficient genetic material that allows the recipient to produce a therapeutically effective amount of Factor VIII so as to reduce, stop and/or prevent hemorrhage. The compositions may be administered alone or in combination with at least one other agent, such as stabilizing compound, which may be administered in any sterile, biocompatible  
20 pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, clotting factors or factor precursors, drugs or hormones. In preferred embodiments, the pharmaceutical compositions also contain a pharmaceutically acceptable excipient. Such excipients include any pharmaceutical  
25 agent that does not itself induce an immune response harmful to the individual receiving the composition, and which may be administered without undue toxicity. Pharmaceutically acceptable excipients include, but are not limited to, liquids such as water, saline, glycerol, sugars and ethanol. Pharmaceutically acceptable salts can be included therein, for example, mineral acid salts such as hydrochlorides,  
30 hydrobromides, phosphates, sulfates, and the like; and the salts of organic acids such

as acetates, propionates, malonates, benzoates, and the like. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles. A thorough discussion of pharmaceutically acceptable excipients is available in *Remington's Pharmaceutical Sciences* (Mack Pub.

5 Co., N.J. [1991]).

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran.

10 Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

20 The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art (e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes).

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 30 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment. For administration of Factor VIII-containing vectors, such labeling would include amount, frequency, and method of administration.

5       Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. Determining a therapeutic effective dose is well within the capability of those skilled in the art using the techniques taught in the present invention, such as ELISA and ChromZ FVIII coagulation activity assay, and other  
10       techniques known in the art. Therapeutic doses will depend on, among other factors, the age and general condition of the subject, the severity of hemophilia, and the strength of the control sequences. Thus, a therapeutically effective amount in humans will fall in a relatively broad range that can be determined through clinical trials.

It is intended that the dosage treatment and regimen used with the present  
15       invention will vary, depending upon the subject and the preparation to be used. Thus, the dosage treatment may be a single dose schedule or a multiple dose schedule. Moreover, the subject may be administered as many doses as appropriate to achieve or maintain the desired blood clotting time.

Direct delivery of the pharmaceutical compositions *in vivo* will generally be  
20       accomplished via injection using a conventional syringe, although other delivery methods such as convention-enhanced delivery are envisioned (*See e.g.*, United States Patent No. 5,720,720, incorporated herein by reference). In this regard, the compositions can be delivered subcutaneously, epidermally, intradermally, intrathecally, intraorbitally, intramucosally (*e.g.*, nasally, rectally and vaginally),  
25       intraperitoneally, intravenously, intraarterially, orally, or intramuscularly. Other modes of administration include oral and pulmonary administration, suppositories, and transdermal applications. In particularly preferred embodiments, the compositions are administered intravenously in the portal vasculature or hepatic artery

One skilled in the art will recognize that the methods and compositions  
30       described above are also applicable to a range of other treatment regimens known in

the art. For example, the methods and compositions of the present invention are compatible with *ex vivo* therapy (e.g., where cells are removed from the body, incubated with the AAV vector and the treated cells are returned to the body).

#### IV. ADMINISTRATION

AAV vector can be administered to any tissue suitable for the expression of Factor VIII. In a preferred embodiments, the AAV vectors of the present invention are successfully administered via the portal vasculature or hepatic artery where it is thought, without being bound by theory, that the vector transduces hepatocytes.

Current approaches to targeting genes to the liver have focused upon *ex vivo* gene therapy. *Ex vivo* liver-directed gene therapy involves the surgical removal of liver cells, transduction of the liver cells *in vitro* (e.g., infection of the explanted cells with recombinant retroviral vectors) followed by injection of the genetically modified liver cells into the liver or spleen of the patient. A serious drawback for *ex vivo* gene therapy of the liver is the fact that hepatocytes cannot be maintained and expanded in culture. Therefore, the success of *ex vivo* liver-directed gene therapy depends upon the ability to efficiently and stably engraft the genetically modified (i.e., transduced) hepatocytes and their progeny. It has been reported that even under optimal conditions, autologous modified liver cells injected into the liver or spleen which engraft represent only a small percentage (less than 10%) of the total number of cells in the liver. Ectopic engraftment of transduced primary hepatocytes into the peritoneal cavity has been tried, in order to address the problem of engraftment in the liver.

Given the problems associated with *ex vivo* liver-directed gene therapy, *in vivo* approaches have been investigated for the transfer of genes into hepatocytes, including the use of recombinant retroviruses, recombinant adenoviruses, liposomes and molecular conjugates. While these *in vivo* approaches do not suffer from the drawbacks associated with *ex vivo* liver-directed gene therapy, they do not provide a means to specifically target hepatocytes. In addition, several of these approaches require performance of a partial hepatectomy, in order to achieve prolonged expression

of the transferred genes *in vivo*. Adenovirus and molecular conjugate based delivery methods also result in liver toxicity and inflammation which is an undesirable feature of Factor VIII gene therapy. The present invention provides compositions and methods for the long-term expression of biologically active Factor VIII. It is contemplated that the present invention will bypass the need for partial hepatectomy, while allowing expression of Factor VIII in concentrations that are therapeutic *in vivo*. The present invention further provides gene therapy compositions and methods that target hepatocytes for the production of Factor VIII by treated individuals.

Other tissues, however, may be suitable for the expression of Factor VIII even if they are not the tissue that normally synthesizes the protein. Muscle cells, for example, have been shown to express biologically active blood clotting Factor IX even though it is normally synthesized in the liver.

Finally, the AAV vectors may contain any nucleic acid sequences coding for biologically active Factor VIII. Additionally, the AAV vectors may contain a nucleic acid coding for fragments of Factor VIII which is itself not biologically active, yet when administered into the subject improves or restores the blood clotting time. For example, as discussed above, the Factor VIII protein comprises two polypeptide chains: a heavy chain and a light chain separated by a B-domain which is cleaved during processing. As demonstrated by the present invention, co-transducing recipient cells with the Factor VIII heavy and light chains leads to the expression of biologically active Factor VIII. Because, however, most hemophiliacs contain a mutation or deletion in only one of the chains (*e.g.*, heavy or light chain), it may be possible to administer only the chain defective in the patient and allow the patient to supply the other chain. In this case, the AAV vector would fall within the scope of the invention even though the single chain (*i.e.*, heavy or light) would not be biologically active until it was administered into a subject which can supply the second chain, thus forming biologically active Factor VIII.

## V. FACTOR VIII ASSAYS

As described in the Experimental section below, there are many ways to assay Factor VIII expression and activity. Although the present invention is not limited to immunoassay methods, the present invention also provides methods for detecting Factor VIII expression comprising the steps of: a) providing a sample suspected of containing Factor VIII, and a control containing a known amount of known Factor VIII; and b) comparing the test sample with the known control, to determine the relative concentration of Factor VIII in the sample. Thus, the methods are capable of identifying samples (*e.g.*, patient samples) with sufficient or insufficient quantities of Factor VIII. In addition, the methods may be conducted using any suitable means to determine the relative concentration of Factor VIII in the test and control samples, including but not limited to means selected from the group consisting of Western blot analysis, Northern blot analysis, Southern blot analysis, denaturing polyacrylamide gel electrophoresis (*e.g.*, SDS-PAGE), reverse transcriptase-coupled polymerase chain reaction (RT-PCR), enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescent immunoassay (IFA). Thus, the methods may be conducted to determine the presence of normal Factor VIII sequences in the genome of the animal source of the test sample, or the expression of Factor VIII (mRNA or protein), as well as detect the presence of abnormal or mutated Factor VIII gene sequences in the test samples.

In one preferred embodiment, the presence of Factor VIII is detected by immunochemical analysis. For example, the immunochemical analysis can comprise detecting binding of an antibody specific for an epitope of Factor VIII. In one another preferred embodiment of the method, the antibody comprises polyclonal antibodies, while in another preferred embodiment, the antibody comprises monoclonal antibodies.

It is further contemplated that antibodies directed against at least a portion of Factor VIII will be used in methods known in the art relating to the localization and structure of Factor VIII (*e.g.*, for Western blotting), measuring levels thereof in appropriate biological samples, etc. The antibodies can be used to detect Factor VIII in a biological sample from an individual (*e.g.*, an individual treated using the methods

WO 00/25110 PC17039/24453

and/or compositions of the present invention). The biological sample can be a biological fluid, including, but not limited to, blood, serum, plasma, interstitial fluid, urine, cerebrospinal fluid, synovial fluid, and the like. In particular, the antigen can be detected from cellular sources, including, but not limited to, hepatocytes. For  
5 example, cells can be obtained from an individual and lysed (*e.g.*, by freeze-thaw cycling, or treatment with a mild cytolytic detergent including, but not limited to, TRITON X-100, digitonin, NONIDET P (NP)-40, saponin, and the like, or combinations thereof; *See, e.g.*, International Patent Publication WO 92/08981).

The biological samples can then be tested directly for the presence of the  
10 Factor VIII using an appropriate strategy (*e.g.*, ELISA or RIA) and format (*e.g.*, microwells, dipstick [*e.g.*, as described in International Patent Publication WO 93/03367], etc.). Alternatively, proteins in the sample can be size separated (*e.g.*, by polyacrylamide gel electrophoresis (PAGE), with or without sodium dodecyl sulfate (SDS), and the presence of Factor VIII detected by immunoblotting [*e.g.*, Western  
15 blotting]). Immunoblotting techniques are generally more effective with antibodies generated against a peptide corresponding to an epitope of a protein, and hence, are particularly suited to the present invention. In another preferred embodiment, the level of Factor VIII is assayed using the whole-blood clotting time and activated partial thromboplastin time (aPTT) of the subject's blood using techniques well known in the  
20 art (Herzog *et al.*, Nature Medicine 5:56 [1999]).

The foregoing explanations of particular assay systems are presented herein for purposes of illustration only, in fulfillment of the duty to present an enabling disclosure of the invention. It is to be understood that the present invention contemplates a variety of immunochemical assay protocols within its spirit and scope.  
25 Indeed, other methods such as biological assays to determine the presence and activity of Factor VIII are also encompassed by the present invention.

Thus, in addition to the immunoassay systems described above, other assay systems, such as those designed to measure and/or detect Fraction VIII and/or clotting

ability of a subject's blood are also encompassed by the present invention (e.g., the ChromZ FVIII coagulation activity [FVIII-c] assay [Helena Labs]).

## EXPERIMENTAL

Below are examples of specific embodiments for carrying out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation should, of course, be allowed for.

In the experimental disclosure which follows, the following abbreviations apply: N (Normal); M (Molar); mM (millimolar);  $\mu$ M (micromolar); g (grams); mg (milligrams);  $\mu$ g (micrograms); ng (nanograms); l or L (liters); ml (milliliters);  $\mu$ l (microliters); cm (centimeters); mm (millimeters);  $\mu$ m (micrometers); nm (nanometers); mU (milliunits);  $^{51}\text{Cr}$  (Chromium 51);  $\mu\text{Ci}$  (microcurie); EC (degrees Centigrade); hFVIII (human factor VIII); FVIII (factor VIII); pH (hydrogen ion concentration); JRH grade; NaCl (sodium chloride); HCl (hydrochloric acid); OD (optical density); bp (base pair(s)); ATP (adenosine 5'-triphosphate); PCR (polymerase chain reaction); DNA (deoxyribonucleic acid); cDNA (complementary DNA); AAV (adeno-associated virus); rAAV (recombinant adeno-associated virus); ITR (inverted terminal repeat); FCS or FBS (fetal calf serum; fetal bovine serum); CFA (complete Freund's adjuvant); BSA (bovine serum albumin); ATCC (American Type Culture Collection, Rockville, MD); Sigma (Sigma Aldrich, St. Louis, MO); Biodesign International (Biodesign International, Kennebunk, ME); Baxter Hyland (Baxter Healthcare Corp., Biotech Group--Hyland Division, Hayward, CA); Helena Labs (Helena Laboratories, Beaumont, TX); American Diagnostica (American Diagnostica, Greenwich, CT); Accurate Chemical (Accurate Chemical and Scientific Corp., Westbury, NY); Molecular Probes (Molecular Probes, Eugene, OR); Vysis (Vysis, Downer Grove, IL); Tel-Test (Tel-Test, Inc., Friendswood, TX); Molecular Dynamics (Molecular Dynamics, Sunnyvale, CA); NUNC (Naperville, IL); and Stratagene

(Stratagene Cloning Systems, La Jolla, CA); and Biodesign (Biodesign International, Kennebunkport, ME).

## EXAMPLE 1

### Dual Vector Plasmid Construction

The heavy and light chains of human Factor VIII (hFVIII) were assembled according to those reported by Yonemura *et al* (Yonemura *et al.*, Prot. Engineer., 6:669-674 [1993]) and cloned as expression cassettes into AAV vectors. Both vectors contain the promoter and the first non-coding intron (from -573 to +985) from the human elongation factor 1 $\alpha$  (EF1 $\alpha$ ) gene (Uetsuki *et al*, J. Biol. Chem., 264:5791-5798 [1989]; and Kim *et al.*, Gene 9:217-223 [1990]). Each vector also contains the first 57 base pairs of the FVIII heavy chain encoding the 19 amino acid signal sequence. The heavy chain construct encodes the A1 and A2 domains and 5 amino acids from the N terminus of the B domain. The light chain vector encodes 85 amino acids of the carboxy terminal B domain, in addition to the A3, C1, and C2 domains. Both vectors utilize the human growth hormone (hGH) polyadenylation signal. The expression cassettes were inserted between AAV ITRs. The initial cloning step involved deleting 854 bp of EF1 $\alpha$  sequences between the *SpeI* and *XcmI* sites of pVm4.1e-hFIX (Nakai *et al.*, Blood 91:1-9 [1998]), and religating to create pVm4.1e $\delta$ D-hFIX.

This construct was then digested with *EcoRI*, which released the hFIX cDNA, and was ligated to an oligonucleotide containing *MfeI* ends (*EcoRI*-compatible) and an internal *ClaI* restriction site, creating pVm4.1e $\delta$ D-linker. The heavy and light chain fragments, including the hGH polyadenylation sequences were isolated from pVm4.1cFVIII-HC and pVm4.1cFVIII-LC, respectively as *ClaI*-*BstEII* fragments. These fragments were cloned between the corresponding sites in the pVm4.1e $\delta$ D-linker, creating plasmids pVm4.1e $\delta$ D-FVIII-HC (also, rAAV-hFVIII-HC) and pVm4.1e $\delta$ D-FVIII-LC (also, rAAV-hFVIII-LC).

Figure 7 provides a map of the constructs. In this figure, the upper line in each panel represents the gene structure of the vectors, and the lower line represents the structure of the hFVIII protein domains encoded by the vectors (ITR, AAV inverted terminal repeat; EF1 $\alpha$  Pro/Intron 1, human polypeptide elongation factor 1  $\alpha$  gene promoter and first intron; hFVIII-HC human FVIII cDNA; hFVIII-LC, human FVIII cDNA; hGH PA, human growth hormone polyadenylation signal; SS, human FVIII signal sequence; A1, A2, "B", A3, C1, C2, complete and incomplete ("") protein domains of the hFVIII protein).

## EXAMPLE 2

### Single Vector Plasmid Construction

The plasmid pAAV-F8-1 construct containing both the light and heavy chains of factor VIII was constructed as follows. A PCR fragment, Z8, containing cloning sites, 5' -splicing donor site of a synthetic intron based on EF1 $\alpha$  and immunoglobulin G (IgG) intron sequences, Kozak sequence and the first 16 nucleotides of the human blood coagulation factor VIII (FVIII) coding sequence was generated using oligonucleotides Z8S and Z8A. The sequences of the nucleic acids is shown below:

Oligonucleotide Z8S:

5' cccaagcttgccggccgcccgggtgccgcccctaggcaggttaagtccgtgtgtggttcc 3'  
(SEQ ID NO:1)

Oligonucleotide Z8A:

5' ccgctcgagcagagctctatttgcattggtggaatcgatgccgcggaaccacacacggc 3'  
(SEQ ID NO:2)

PCR fragment Z8:

5' cccaagcttgccggccgcccgggtgccgcccctaggcaggttaagtccgtgtgtggttccgc  
ggcatcgattccaccatgcaaatagagctctgctcgagcgg 3' (SEQ ID NO:3)

Nucleic acid Z8 was inserted into pZERO-2 (Invitrogen) between *HindIII* and *XhoI* sites to create pZ8. A PCR fragment, INT3, containing the branching point, the

polypyrimidine tract, and the 3' splicing acceptor site of the synthetic intron was generated using oligonucleotides INT3S and INT3A whose sequence is shown below.

Oligonucleotide INT3S:

5            5' ttccgcgggcctggcctctttacgggttatggcccttgcgtgccttgaattactga 3'  
             (SEQ ID NO:4)

Oligonucleotide INT3A:

             5' gaatcgatacctgtggagaaaaagaaaaagtggatgacagtgcagtaattcaaggc 3'  
10            (SEQ ID NO:5)

PCR Fragment INT3:

             5' ttccgcgggcctggcctctttacgggttatggcccttgcgtgccttgaattactgacact  
             gacatccactttttctttctccacaggtatcgattc 3' (SEQ ID NO:6)  
15

INT3 was inserted between the *Sac*II and *Cla*I sites of pZ8 to create pZ8.I. Therefore, Z8.I contains the entire synthetic intron between *Avr*II and *Cla*I sites. A hFVIII cDNA fragment having *Sac*I and *Xho*I restriction sites was inserted between the *Sac*I and *Xho*I sites of pZ8.I to create pZ8.I.dB. Therefore, pZ8.I.dB contains a  
20 synthetic intron and the entire coding sequence of hFVIII.

A PCR fragment, EG3, containing three HNF-3 binding sites and -54 to +8 of mouse albumin gene was generated using oligonucleotides EG3S and EG3A with modifications to eliminate linker sequences. The sequences of EG3S and EG3A are as follows:

25

Oligonucleotide EG3S:

             5' agggaatgtttgttcttaaataccatccaggggaatgtttgttcttaaataccatccaggggaatgtttgttcttaaatacca  
             tctacagttattgggtaaa 3' (SEQ ID NO:7)  
30

Oligonucleotide EG3A:

5' ggaaaggatgatctgtgtgcagaaagactcgctctaataacttctttaaccaataactg 3'  
(SEQ ID NO:8)

5 PCR Fragment EG3:

5' aggggaatgtttgttcttaaataccatccaggggaatgtttgttcttaaataccatccaggggaatgtttgttctta  
aataccatctacagtattggttaaagaagtattagagcgagctcttctgcacacagatcacctttcc 3'

(SEQ ID NO:9)

10 EG3 was then phosphorylated using T4 polynucleotide kinase and inserted into the *Sma*I site of pZ8.I.dB to create pZ8.I.dB.egg. A DNA fragment, SPA, containing an efficient synthetic polyA signal based on rabbit  $\beta$ -globin sequences (Genes and Develop., 3:1019) was generated by hybridizing two oligonucleotides SPA.S and SPA.A.

15

Oligonucleotide SPA.S:

5' tcgagaataaaagatcagagctctagagatctgtgtgttggtttttgtgtgcggccgc 3'  
(SEQ ID NO:10)

20 Oligonucleotide SPA.A:

5' tcgagcggccgcacacaaaaaccaacacacagatctctagagctctgatcttttattc 3'  
(SEQ ID NO:11)

PCR Fragment SPA:

25 5' tcgagaataaaagatcagagctctagagatctgtgtgttggtttttgtgtgcggccgctcga 3'  
(SEQ ID NO:12)

SPA was inserted into the *Xho*I site of pZero-2 to create pZero-2.SPA. SPA was excised from a pZero-2.SPA clone and inserted into the *Xho*I site of pZ8.I.dB.egg

to create pZ8.I.dB.egg.A. pAAV-CMV-FIX9 was digested with *Cla*I, blunted with T4 polymerase and religated to create pAAV(*Cla*-)-CMV-FIX9.

The entire expression cassette containing HNF-3.albumin promoter-synthetic intron-hFVIII-synthetic poly A signal was excised from pZ8.I.dB.egg.A using *Not*I and  
5 ligated to the plasmid backbone and AAV ITRs from pAAV (*Cla*-)-CMV-FIX9 to create pAAV-F8-1. The nucleotide sequence of the vector from ITR to ITR (*i.e.*, excluding plasmid backbone) is shown in SEQ ID NO 13.

### EXAMPLE 3

#### Virion Production

AAV vectors were produced from these plasmids using the Ad free system as previously described in U.S. Patent No. 5,858,351; U.S. Patent No. 5,846,528; U.S. Patent No. 5,622,856; and Matsushita *et al.*, Gene Ther 5:938 (1998) all of which are hereby incorporated by reference. Briefly, 293 cells (ATCC, catalog number CRL-  
15 1573) were seeded in 10 cm dishes at a density of  $3 \times 10^6$  cells per dish in 10 ml medium and incubated at 37°C with CO<sub>2</sub> and humidity. After an overnight incubation, cells were approximately seventy to eighty percent confluent.

The cells were then transfected with DNA by the calcium phosphate method, as is well known in the art. Briefly, 7 µg of AAV vector containing the Factor VIII  
20 coding region, 7 µg of pladen05 which supplies the accessory functions, and 7 µg of 1909 AAV helper were added to a 3 ml sterile, polystyrene snap cap tube using sterile pipette tips. Then, 1.0 ml of 300 mM CaCl<sub>2</sub> (JRH grade) was added to each tube and mixed by pipetting up and down. An equal volume of 2X HBS (274 mM NaCl, 10 mM KCl, 42 mM HEPES, 1.4 mM Na<sub>2</sub>PO<sub>4</sub>, 12 mM dextrose, pH 7.05, JRH grade)  
25 was added with a 2 ml pipette, and the solution was pipetted up and down three times. The DNA mixture was immediately added to the cells, one drop at a time, evenly throughout the 10 cm dish. The cells were then incubated at 37°C with CO<sub>2</sub> and humidity for six hours. A granular precipitate was visible in the transfected cell cultures. After six hours, the DNA mixture was removed from the cells, which were  
30 provided with fresh medium and incubated for 72 hours.

After 72 hours, the cells were harvested, pelleted, and resuspended in 1 ml TBS/1% BSA. Freeze/thaw extracts were prepared by repeatedly (three times) freezing the cell suspension on dry ice and thawing at 37°C. Viral preps were stored at -80°C and titered by dot blot assay prior to the first round of infection.

5

#### EXAMPLE 4

##### *In Vitro* Cell Transduction

Cells from the stable human cell line, 293 (ATCC No. CRL1573) were seeded in six-well plates (*i.e.*, plates having six wells for cell growth) at a density of  $5 \times 10^5$  cells/well. When the monolayers reached 80-90% confluence, they were infected with rAAV virions AAV-e8D-FVIII-HC, AAV-e8D-FVIII-LC, an equal ratio of AAV-e8D-FVIII-HC and AAV-e8D-FVIII-LC, or AAV-e8D-FIX at MOIs of  $3 \times 10^3$  and  $3 \times 10^4$ . Eighteen hours post infection, the media were replaced with DMEM/10% heat inactivated FBS. The media were collected later for analysis by ELISA (as described below) for FVIII light chain antigen levels, and by the ChromZ FVIII as coagulation activity (FVIII-c) assay (Helena Labs) for biological activity, using the manufacturer's instructions and as described in Example 6.

10

15

#### EXAMPLE 5

##### Single Chain Factor VIII Infectivity Assay

20

25

30

In this Example, the infectivity of single chain Factor VIII was investigated. To determine the infectivity of rAAV-hF8-1, HepG2, 293, and H2.35 cells were infected with rAAV-hF8-1 and a control vector rAAV-hF8L at an MOI of  $1 \times 10^4$  viral particles per cell. Recombinant AAV DNA in infected cells was isolated by Hirt extraction and run on an alkaline agarose gel. Southern blot analysis using an human F8 probe showed that similar amounts of rAAV-hF8-1 and rAAV-hF8L were isolated from uncoated virus in the infected cells. An infectious center assay (ICA) known in the art (*See e.g.*, Snyder, *Current Protocols in Genetics*, Chapter 12, John Wiley & Sons [1997]) was used to further characterize the infectivity of rAAV-hF8-1. In this assay, the infectious particle to total particle ratio of rAAV-F8-1 and that of a control

rAAV vector with the genome size of 4645 nucleotides was determined. The results indicated that rAAV-hF8-1 had an infectious particle to total particle ratio that was comparable to the control vector at approximately 1:1000. Taken together, these results indicate rAAV-hF8-1 has similar infectivity as rAAV vectors with the genome size of wild-type AAV.

## EXAMPLE 6

### Factor VIII Protein Expression Assay

An ELISA specific for the light chain of FVIII was used to determine FVIII light chain antigen levels in the 293 cells, as well as the injected animals (described below). NUNC Maxisorb 96 well plates were coated with 50  $\mu$ l of a 1:500 dilution of the light chain specific antibody, N77110 (Biodesign International) in a coating buffer overnight at 4°C. The plate was washed three times with wash buffer (PBS, 0.05% Tween 20) and blocked with 200  $\mu$ l blocking buffer (PBS, 10% horse serum, 0.05% Tween 20) at room temperature for 1 hour. The plate was washed three times and standards and samples were applied. Bioclone recombinant human FVIII (Baxter Hyland) was used as the standard, and was diluted in blocking buffer to concentrations ranging from 320 ng/ml to 10 ng/ml.

For analysis of transduced culture supernatants, the standards contained 50% media, and for analysis of mouse plasma, the standards were diluted into 10%, in normal pooled mouse plasma (Sigma). A standard assay reference plasma (SARP; Helena Labs) was also included in the assay. Following the loading of the standards and samples (95  $\mu$ l/well), the plate was incubated at room temperature for 2 hours, and washed five times with wash buffer (200  $\mu$ l/well). A 1:200 dilution of a horseradish peroxidase-conjugated light chain specific antibody, ESH8-HRP, (American Diagnostica) was added (100  $\mu$ l/well), and the plate was incubated for 1 hour at room temperature. The plates were then washed four times with wash buffer, and the antigen was detected using an ABTS peroxidase substrate kit (BioRad) according to the manufacturer's instructions. The results are shown in Table 1 of Example 7, below.

## EXAMPLE 7

### Factor VIII Biological Activity Assay

The ChromZ FVIII coagulation activity (FVIII-c) assay (Helena Labs, Beaumont, TX) was used to detect biologically active FVIII in the 293 cells infected as described in Example 4. Bioclone recombinant human FVIII (Baxter Hyland) was used as a standard to analyze transfected culture supernatants. The standards were diluted in plasma dilution buffer (supplied in kit) in the range of 10 ng/ml to 0.313 ng/ml, and were made 2.5% in media. Because this assay can detect both human and murine FVIII activity, it was modified to deplete biologically active human Factor VIII in the mouse plasma. Mouse plasma was pre-incubated with an antibody specific for human FVIII prior to performing the assay. The difference in FVIII activity between the untreated plasma sample and the antibody treated sample represent the amount of biologically active human FVIII in the plasma. The standard used in the assay was normal pooled human plasma (FACT; obtained from George King Biomedical). Serial dilutions of FACT were made in FVIII deficient plasma from undiluted (200 ng/ml) to 6.25 ng/ml. The standards (10  $\mu$ l) were incubated at 37°C for 15 min., with or without the addition of 2  $\mu$ l antibody N77110. Similarly, mouse plasma samples were diluted in FVIII deficient plasma and 10  $\mu$ l of these diluted samples were incubated with or without 2  $\mu$ l of N77110 at 37°C for 15 min., and immediately placed on ice. Thus, all incubations with antibody were done in a background of 100% plasma. The antibody adsorbed and non-adsorbed FACT standards, as well as the mouse plasma samples were diluted 1:20 in plasma detection buffer provided in the ChromZ kit. Thus, the final concentration of the FACT standards used in the assay ranged from 10 ng/ml to 0.313 ng/ml.

Twenty five microliters of these dilutions were added to a chilled 96 well plate. With the plate on ice, 25  $\mu$ l of FIXa reagent and 50  $\mu$ l of FX were added, and the plate was incubated at 37°C for 15 min. Substrate (50  $\mu$ l) was added and the plate was incubated for an additional 3 min at 37°C. The reaction was stopped with the addition of 25  $\mu$ l 50% acetic acid and the optical density at 405 nm was measured.

biologically active FVIII. It was assumed that equal amounts of heavy and light chains were produced in transduced cells. The activity units were converted to nanograms using the definition of one unit being equal to the amount of FVIII in 1 ml of normal pooled human plasma, or 200 ng.

**TABLE 1. *In Vitro* Production of Biologically Active  
Human Factor VIII From Two rAAV Vectors**

Vector	MOI	ELISA (ng/ml)	ChromZ	
			(mU/ml)	(ng/ml)
AAV-e8D-FVIII-HC and AAV-e8D-FVIII-LC	$3 \times 10^3$	24	35	7.1
AAV-e8D-FVIII-HC and AAV-e8D-FVIII-LC	$3 \times 10^4$	121	440	87.9
AAV-e8D-FVIII-HC	$3 \times 10^3$	0	0	0
AAV-e8D-FVIII-HC	$3 \times 10^4$	0	0	0
AAV-e8D-FVIII-LC	$3 \times 10^3$	20.5	0	0
AAV-e8D-FVIII-LC	$3 \times 10^4$	96.9	0	0
AAV-hFIX	$3 \times 10^3$	0	0	0
AAV-hFIX	$3 \times 10^4$	0	0	0
No Vector		0	0	0

### EXAMPLE 8

#### Immunofluorescent Staining of FVIII Heavy and Light Chains

In these experiments, 293 cells transduced as described above were analyzed using immunofluorescent staining. 293 cells were plated on rat tail collagen-coated two-well culture slides at a density of  $4 \times 10^5$  cells per well. Forty-eight hours later, the cells were transduced at an MOI of  $3 \times 10^4$  particles per cell of rAAV-hFVIII-HC and rAAV-hFVIII-LC. Forty-eight hours post-transduction, the cells were fixed *in situ* with acetone, blocked with 2% BSA, and stained with a fluorescently labelled anti-hFVIII light chain antibody and a fluorescently labelled anti-hFVIII heavy chain antibody. The anti-hFVIII light chain antibody used was ESH-4 monoclonal antibody (American Diagnostica), fluorescently labelled with alexa-488 (Molecular Probes),

according to the manufacturer's instructions. The anti-hFVIII heavy chain antibody used was MAS530P monoclonal antibody (Accurate Chemical) fluorescently labelled with alexa-594 (Molecular Probes), according to the manufacturer's instructions. The cells were counter-stained with DAPI. The images were collected using a Zeiss  
5 Axioskop fluorescence microscope equipped with separate filters for DAPI, FITC, and rhodamine signals and a CCD camera. Image analysis was performed using Quips imaging software (Vysis).

As indicated above, infection of cells with either rAAV-e8D-FVIII-HC or AAV-e8D-FVIII-LC, followed by staining with antibodies to both chains resulted in  
10 production of the individual chains of human FVIII. Immunofluorescent staining of cells co-infected with both vectors demonstrated that although some cells express only the heavy or light chain of hFVIII, many co-expressed both chains of human FVIII.

## EXAMPLE 9

### *In Vitro* Expression of Factor VIII Using Single Construct

Table 2 shows that two single vector constructs containing the heavy and the light chain of Factor VIII driven by different promoters express biologically active Factor VIII. The constructs pAAV-hF8-1 (SEQ ID NO:13), and pVm4.1cF8ΔB (SEQ ID NO:14) were transfected into 293 cells. Following transfection, the cells were  
20 allowed to express factor VIII for 48-72 hours. Factor VIII in the culture media was assayed by the ChromZ FVIII coagulation activity (FVIII-c) assay, as per the manufacturer's instructions.

TABLE 2

*In Vitro* Production of Biologically Active FVIII

Construct(s)	ELISA (ng/ml)	ChromZ (ng/ml)
Control	-	0
pAAV-hF8-1	-	4.9
pVm4.1cF8ΔB	-	46

## EXAMPLE 10

## Factor VIII Expression Using Tissue Specific Promoters

In these experiments, different promoters and enhancer elements were used to drive expression of a Factor VII coding sequence. Expression of Factor VIII was compared in 293 cells and HepG2 cells using different promoters. The pAAVeF8ΔB contains an EF-1 $\alpha$  promoter with a hGH intron, Factor VIII with a B-domain deletion (F8ΔB) and a polyA. As described previously, pAAV-hF8-1 uses the HNF-3 albumin promoter with a minimal intron followed by F8ΔB and a minimal polyA. The construct pAAV-c8 uses the CMV enhancer-promoter and the F8ΔB. pAAV8b1 contains the HNF-3 albumin promoter followed by the CMV/B-globin intron with the F8ΔB and a minimal poly A site. Table 3 describes Factor VIII expression using the albumin promoter relative to the control plasmid pV4.1cF8ΔB in HepG2 and 293 cells. These data show increased expression of Factor VIII in HepG2 liver cells with the albumin promoter as compared to Factor VIII expression in 293 cells.

TABLE 3

## Relative Tissue Specificity of Promoters

Plasmid Construct	HepG2 Cells	293 Cells
pAAV-hF8-1	6.2	0.6
pAAV8bl	6.7	1.0
pAAVc8	30.0	41.0
pV4.1eF8ΔB	100	100

Next, several promoters derived from the transthyretin (TTR) gene promoter were transfected into HepG2 cells. TTR is an abundant serum protein and the gene enhancer-promoter contains well known liver-specific transcription factor binding sites (Samadani *et al.*, Gene Expression 6:23 [1996]; Yan *et al.*, EMBO 9:869 [1990]; Costa and Grayson, Nuc. Acids Res., 19:4139 [1991]; Costa *et al.*, Mol. Cell. Bio., 6:4697 [1986]). The constructs were made by replacing the HNF-3 albumin promoter in pAAV-hF8-1 with various lengths of the TTR promoter-enhancer. The TTR enhancer-promoter was modified by replacing the weak affinity binding sites with the strong affinity binding sites to create pAAV-hF8-2. The pAAV-hF8-TTR-E-L-P202 construct contains the full TTR promoter with a linker between the enhancer and the promoter. The remaining constructs are 5' deletions: pAAV-hF8-TTR-E-P202 has the promoter and enhancer with no linker; pAAV-hF8-TTR-E-P197 has a 5 base pair deletion from the promoter; pAAV-hF8-TTR-E-P151 has a 50 base pair deletion; pAAV-hF8-TTR-P202 lacks the TTR enhancer and pAAV-hF8-TTR(X) has a 65 base pair deletion in the enhancer. The control plasmid, pAAV-hF8-1, expressed approximately 4.6 mU/ml. Table 4 shows the fold-increase in Factor VIII activity using the TTR promoter series relative to the control plasmid.

TABLE 4

## Factor VIII Expression Using TTR-Derived Promoters

Plasmid Construct	Relative Factor VIII Activity
pAAV-hF8-STTR	3.16
pAAV-hF8-TTR-E-L-P202	8.86
pAAV-hF8-TTR-E-P202	6.1
pAAV-hF8-TTR-EP197	7.3
pAAV-hF8-TTR-E-P151	13.3
pAAV-hF8-TTR-P202	2.3

## EXAMPLE 11

*In Vivo* Expression of Factor VIII

In order to test the feasibility of the AAV vector approach of the present invention *in vivo*, three groups of five C57BL/6 mice were injected via the portal vein with either  $3 \times 10^{11}$  particles of AAV-e $\delta$ D-FVIII-HC,  $3 \times 10^{11}$  particles of AAV-e $\delta$ D-FVIII-LC, or  $3 \times 10^{11}$  particles of both AAV-e $\delta$ D-FVIII-HC and AAV-e $\delta$ D-FVIII-LC. In addition, a group of four animals was injected with  $3 \times 10^{11}$  particles of AAV-e $\delta$ D-FIX. It has been shown that this strain of mice does not elicit an immune response to human FVIII when the gene is delivered to the portal vein via an adenoviral vector (Connelly *et al.*, Blood 87:4671-4677 [1996]). As indicated by the results shown below, the data obtained during these experiments demonstrate the feasibility of producing biologically active FVIII using two AAV vectors to independently deliver the heavy and light chains of FVIII.

Blood samples were collected in sodium citrate via the retro-orbital plexus at biweekly intervals for the first 2 months and at monthly intervals thereafter for 6

FVIII remaining in the plasma after adsorption represented the amount of active murine FVIII and the difference represented the amount of active human FVIII. Control experiments demonstrated that the antibody could remove 80-90% of the human FVIII from a mouse plasma sample when the sample was spiked with up to 32 ng of human FVIII. The modified ChromZ assay indicated that only those animals injected with both vectors produced biologically active FVIII, as indicated in Table 5. The results shown in Table 5 are those from plasma collected 8 weeks post-injection, although similar results were obtained at 10 weeks and 5 months post-injection. One of the five animals co-injected with both the heavy and light chain vector did not express VIII, presumably due to an inefficient injection, and was omitted from the analysis. Animals injected with both vectors produced over 2 µg/ml hFVIII light chain as measured by ELISA. The ChromZ assay indicated that a total of 600-900 ng/ml of active hFVIII was detected in the plasma. The contribution from murine Factor VIII was approximately 400-500 ng/ml, indicating that about 230-430 ng/ml of active human Factor VIII was present in the plasma. Although only a fraction of the total protein was found to be active, the animals produced physiological levels of the active protein (*i.e.*, 200 ng/ml). The animals were found to have maintained these physiological levels of active protein for more than 11 months, without waning. Similar analyses performed on animals injected with the light chain vector alone, the heavy chain vector alone, or the hFIX vector demonstrated no biologically active human FVIII in the plasma of these animals.

TABLE 5

Biological Activity of Human Factor VIII *In Vivo*

Construct(s) Used	ELISA (ng/ml)	Total FVIII (-Ab) (Units)	Murine FVIII (+Ab) (Units)	Human FVIII (ng/ml)
AAV-e8D-FVIII-HC and AAV-e8D-FVIII-LC*	2288	3.9	2.2	342
AAV-e8D-FVIII-LC *	3329	1.4	1.6	0
AAV-e8D-FVIII-HC	0	1.6	1.6	0
AAV-e8D-FIX	0	1.4	2.0	0

\*Average of three animals.

## EXAMPLE 12

## Gene Transfer and Vector Expression in Tissues

In these experiments, evidence of gene transfer to liver was obtained by Southern Blot analysis following isolation of DNA from one animal of each experimental group sacrificed 8 weeks post-injection (*i.e.*, as described in Example 11). In addition, DNA was obtained from other tissues in order to determine the degree of vector expression in organs other than the liver.

Twenty micrograms of DNA was digested with *Bgl*II, separated using a 1% agarose gel, and hybridized with a <sup>32</sup>P-labelled 1126 bp *A*/wNI fragment encoding the A1 and A2 domains of hFVIII (heavy chain probe), or a <sup>32</sup>P-labelled 1456 bp *Nde*I-*Eco*RI fragment encoding the A3, C1 and C2 domains of hFVIII (light chain probe).

Copy number controls were generated by spiking *Bgl*II-digested naive mouse liver DNA with *Bgl*II-digested heavy or light chain plasmid DNA (pVm4.1e8D-hFVIII-HC and pVm4.1e8D-hFVIII-LC, respectively), at ratios of 10, 5, 1, 0.1, and 0.01 copies per diploid genome. The hybridized membranes were analyzed using a Storm 860 phosphorimager (Molecular Dynamics), and quantitation of vector copy number was evaluated using ImageQuaNT software (Molecular Dynamics). Autoradiography of the

hybridized membranes was also performed. Total RNA was isolated from liver tissue using the RNA Stat extraction kit (Tel-Test). As describe briefly below, Northern blot analysis was also performed on 10 µg RNA using methods known in the art, in conjunction with the <sup>32</sup>P-labelled probes specific to the heavy and light chains of hFVIII described above and autoradiography was performed on the hybridized membranes.

Following digestion of liver DNA with *Bgl*III and hybridization with an hFVIII light chain probe described below, using methods known in the art, a band at the predicted size of 3015 bp was detected in animals injected with rAAV-hFVIII-LC, or both the heavy and light chain vectors. This band was not observed in the DNA of animals injected with the heavy chain vector alone or the hFIX vector, as shown in Figure 9A (rAAV-hFVIII-LC, lane 1; rAAV-hFIX, lane 2; rAAV-hFVIII-HC, lane 3; both rAAV-hFVIII-LC and rAAV-hFVIII-HC, lane 4; copy number controls were generated by spiking *Bgl*III digested naive mouse liver DNA with the corresponding plasmids at ratios of 10, 5, 1, 0.1, and 0.01 copies per diploid genome, lanes 5-9). Phosphoimage analysis revealed that the light chain vector was present at approximately 2.4 and 1.5 copies per diploid genome in animals injected with the light chain vector alone or both vectors, respectively. When *Bgl*III-digested DNA was hybridized with an hFVIII heavy chain probe, the expected band of 2318 bp was observed in animals injected with the heavy chain vector alone or both vectors, but was not detected in animals injected with the light chain vector alone or the hFIX vector, as shown in Figure 9B. The copy number in animals injected with the heavy chain vector alone and both vectors was 1.1 and 1.7 vector copies per diploid genome, respectively.

The results of hybridization of DNA extracted from the spleen, kidney and heart tissue with either an hFVIII light chain probe or a heavy chain probe indicated that these tissues contained less than 1 copy of vector sequences per 10 diploid genomes, demonstrating that the vector distributes primarily to the liver following intra-portal injection, as shown in Figures 10A and 10B.

Human FVIII gene expression in the liver of the mice was also assessed by Northern blot analysis on RNA isolated from animals sacrificed 8 weeks post-injection (as described above). hFVIII light chain transcripts of the predicted size (2.7 kb) were observed in animals injected with the light chain vector alone or both vectors, as shown in Figure 11A. Similarly, the expected hFVIII heavy chain transcripts (2.7 kb) were detected in animals that were injected with the heavy chain vector alone or both vectors, as shown in Figure 11B. Since the heavy and light chain DNA sequences were shown by Southern blot analysis to be present at approximately the same copy number (1.7 and 1.5 copies per diploid genome, respectively), in an animal injected with both vectors, these results demonstrate that both the heavy and light chains of hFVIII are expressed in the liver in approximately equivalent amounts.

## CLAIMS

We claim:

1. A method of treating hemophilia in a human comprising:

- 5           a) providing at least one recombinant adeno-associated vector comprising nucleotide sequences encoding Factor VIII; and
- b) administering said at least one recombinant adeno-associated vector into said subject under conditions such that said Factor VIII nucleotide sequences are expressed at a level which
- 10           provides a therapeutic effective amount in said subject.

2. The method of Claim 1, wherein said Factor VIII nucleotide sequences are expressed in the liver of said subject.

15           3. The method of Claim 1, wherein said blood clotting disorder is ameliorated.

          4. The method of Claim 1, wherein said blood clotting disorder is hemophilia A.

20           5. The method of Claim 1, wherein said subject is a human.

          6. The method of Claim 1, wherein said recombinant adeno-associated vector encodes the light chain of said Factor VIII.

25           7. The method of Claim 1, wherein said recombinant adeno-associated vector encodes the heavy chain of said Factor VIII.

8. The method of Claim 1, wherein said Factor VIII nucleotide sequences encode at least one Factor VIII domain selected from the group consisting of A1, A2, A3, B, C1, and C2.

5 9. The method of Claim 1, wherein said recombinant adeno-associated vector is administered intravenously.

10. The method of Claim 8, wherein said intravenous administration is via the portal vein.

10 11. The method of Claim 1, wherein said recombinant adeno-associated vector further comprises nucleotides encoding the human growth hormone polyadenylation sequence.

15 12. A method of treating a subject comprising:

a) providing:

i) a subject suffering from a blood clotting disorder,

ii) a first recombinant adeno-associated vector comprising nucleotide sequences encoding the light chain of Factor VIII, and

20 iii) a second recombinant adeno-associated vector comprising nucleotide sequences encoding the heavy chain of Factor VIII; and

25 b) administering said at first and second recombinant adeno-associated vector into said subject under conditions such that said Factor VIII heavy and light chain nucleotide sequences are expressed at a level which provides a therapeutic effect in said subject.

13. The method of Claim 12, wherein said Factor VIII heavy and light chain nucleotide sequences are expressed in the liver of said subject.

14. The method of Claim 12, wherein said Factor VIII nucleotide sequences  
5 encode at least one Factor VIII domain selected from the group consisting of A1, A2, A3, B, C1, and C2.

15. The method of Claim 12, wherein said blood clotting disorder is ameliorated.

16. The method of Claim 12, wherein said blood clotting disorder is hemophilia A.

17. The method of Claim 12, wherein said subject is a human.

18. The method of Claim 12, wherein said recombinant adeno-associated vector is administered intravenously.

19. The method of Claim 18, wherein said intravenous administration is via  
20 the portal vein.

20. The method of Claim 12, wherein said recombinant adeno-associated vector further comprises nucleotides encoding the human growth hormone polyadenylation sequence.

21. A recombinant adenovirus-associated vector, wherein said vector  
comprises nucleic acid encoding at least a portion of Factor VIII operably linked to

control sequences, wherein said control sequences direct the transcription of translation of said portion of Factor VIII, and a pharmaceutically acceptable excipient.

22. The vector of Claim 21, wherein said nucleic acid encoding Factor VIII  
5 comprises nucleotides encoding the heavy chain of Factor VIII.

23. The vector of Claim 21, wherein said nucleic acid encoding Factor VIII  
comprises nucleotides encoding the light chain of Factor VIII.

10 24. The vector of Claim 21, wherein said nucleic acid encoding at least a  
portion of Factor VIII encodes at least one Factor VIII domain selected from the group  
consisting of A1, A2, A3, B, C1, and C2.

25. The vector of Claim 21, wherein said vector further comprises  
15 nucleotides encoding the human growth hormone polyadenylation sequence.

26. A host cell containing at least one recombinant adeno-associated vector,  
wherein said host cell is capable of expressing Factor VIII.

20 27. The host cell of Claim 26, wherein said cell is a liver cell.

28. The host cell of Claim 26, wherein said Factor VIII is expressed at or  
above physiological levels.

25 29. The host cell of Claim 26, wherein said host cell expresses said Factor  
VIII at a level greater than about 200 ng/ml.

30. The host cell of Claim 29, wherein said host cell is present in an animal.

FIGURE 1




FIGURE 2




FIGURE 3

## AAV-F8-1




FIGURE 4

## PVM4.1c-F8AB




FIGURE 5

CAGCTGCGCGCTCGCTCGCTCACTGAGGCCGCCCGGGCAAAGCCCCGGGCGTCGGGCGACCTTTGGTCGCCCCGGCCTCAGT  
GAGCGAGCGAGCGCGCAGAGAGGGAGTGGCCAACTCCATCACTAGGGGTTCTGCGGCCGCCAGGGAATGTTTGTCTT  
AAATACCATCCAGGGAATGTTTGTCTTAAATACCATCCAGGGAATGTTTGTCTTAAATACCATCTACAGTTATTGGTT  
AAAGAAGTATATTAGAGCGAGTCTTCTGCACACAGATCACTTTCCGGGTGCCGCCCTAGGCAGGTAAGTGCCGTGTG  
TGGTTCCCGCGGGCCTGGCCTCTTTACGGGTTATGGCCCTTGCGTGCCTTGAATTACTGACACTGACATCCACTTTTCT  
TTTTCTCCACAGGTATCGATTCCACCATGCAAATAGAGCTCTCCACCTGCTTCTTCTGTGCTTTTGCGATTCTGCTTT  
AGTGCCACCAGAAGATACTACCTGGGTGCAGTGGAACTGTCTATGGGACTATATGCAAAGTGATCTCGGTGAGCTGCCTGT  
GGACGCAAGATTTCTCTAGAGTGCCAAAATCTTTCCATTCACACCTCAGTCGTGTACAAAAAGACTCTGTTTGTAG  
AATTACGGATCACTTTTCAACATCGCTAAGCCAAGGCCACCTGGATGGGTCTGCTAGGTCTTACCATCCAGGCTGAG  
GTTTATGATACAGTGGTCACTTACACTTAAGAACATGGCTTCCCATCTGTCTGCTCTTCTGCTGTTGGTGTATCCTACTG  
GAAAGCTTCTGAGGGAGCTGAATATGATGATCAGACCAGTCAAAGGGAGAAAGATGATAAAGTCTTCCCTGGTGGAA  
GCCATACATATGTCTGGCAGGTCTGAAAGAGAATGGTCCAATGGCCTCTGACCCACTGTGCCTTACCTACTCATATCTT  
TCTCATGTGGACCTGGTAAAAGACTTGAATTGAGGCTCATTTGGAGCCCTACTAGTATGTAGAGAAGGGAGTCTGGCCAA  
GGAAAAGACACAGACCTTGCACAAATTTATACTACTTTTCTGCTGATTTTGTATGAAGGGAAAAGTTGGCATCAGAAACAA  
AGAACTCCTTGATGCAGGATAGGGATGCTGCATCTGCTCGGGCCTGGCCTAAAATGCACACAGTCAATGGTTATGTAAAC  
AGGTCTCTGCAGGTCTGATTGGATGCCACAGGAAATCAGTCTATTGGCATGTGATTTGGAATGGGCACCCTCTGAAGT  
GCACTCAATATTCTCGAAGGTCAACATTTCTTGTGAGGAACCATCGCCAGGCGTCTTGGAAATCTCGCCAATAACTT  
TCCTTACTGTCTCAACACTCTTGATGGACCTTGGACAGTTTCTACTGTTTGTCTATCTCTTCCCAACATCATGTGGC  
ATGGAAGCTTATGTCAAAGTAGACAGCTGTCCAGAGGAACCCCACTACGAATGAAAAATAATGAAGAAGCGGAAGACTA  
TGATGATGATCTTACTGATTCTGAAATGGATGTGGTCAAGTTTGTATGATGACAACCTCTCTTCTTTATCCAAATTCGCT  
CAGTTGCCAAGAAGCATCTAAAACCTTGGGTACATTACATGTCTGCTGAAGAGGAGGACTGGGACTATGTCCCTTAGTC  
CTCGCCCCGATGACAGAAGTTATAAAAGTCAATATTTGAACAATGGCCCTCAGCGGATTGGTAGGAAGTACAAAAAGT  
CCGATTTATGGCATACACAGATGAAACCTTTAAGACTCGTGAAGCTATTGAGCATGAATCAGGAATCTTGGGACCTTAC  
TTTATGGGGAAGTTGGAGACACACTGTTGATTATATTTAAGATAATCAAGCAAGCAGACCATATAACATCTACCTCAGCGA  
ATCACTGATGTCCGTCTTTGTATTCAAGGAGATTACCAAAAGGTGTAACATTTGAAGGATTTTCCAATCTGCCAGG  
AGAAATATTCAAATATAAATGGACAGTGTAGTGAAGATGGGCCAACTAAATCAGATCCTCGGTGCCTGACCCGCTATT  
ACTCTAGTTTCTGTTAATATGGAGAGAGATCTAGCTTCAGGACTCATTGGCCCTCTCTCATCTGCTACAAAGAATCTGTA  
GATCAAAGAGGAAACCAGATAATGTCAGACAAGAGGAATGTCATCTGTTTCTGTTTGTATTTGATGAGAACCGAAGCTGTA  
CCTCAGAGGAATATACAACGCTTTCTCCCCAATCCAGTTGGAGTCAGCTTGAGGATCCAGAGTTCCAGCCTCCAACA  
TCATGCACAGCATCAATGGCTATGTTTTGTATGATGTTTGCAGTTGTGATGTTTGTGATGAGGTGGCATACTGGTACATT  
CTAAGCATTTGAGCACAGACTGACTTCTTTCTGTCTTCTTCTGATATACCTTCAAACACAAATGGTCTATGAAGA  
CACACTCACCTATTCCCATTTCTCAGGAGAACTGTCTTCATGTGATGGAACCCAGGTCTATGGATTCTGGGGTGCC  
ACAACCTGACTTTCCGAAACAGAGGCATGACCCGCTTACTGAAGGTTTCTAGTTGTGACAAGAACACTTGGTATTATTAC  
GAGGACAGTTATGAAGATATTTTCAGCATACTTGTCTGATGATAAATAATGCCATTGAACCAAGAAGCTTCGAAATAACTCG  
TACTACTCTTCAGTCAGATCAAGAGGAAATTGACTATGATGATACCATATCAGTTGAAATGAAGAAGGAAGATTTTGACA  
TTTATGATGAGGATGAAAATCAGAGCCCCCGCAGCTTTCAAAGAAAAACACGACACTATTTTATTGCTGCAGTGGAGAGG  
CTCTGGGATTATGGGATGAGTAGCTCCCCACATGTTCTAAGAAACAGGGCTCAGAGTGGCAGTGTCCCTCAGTTCAAGAA  
AGTTGTTTTTCAGGAATTTACTGATGGCTCCTTTACTCAGCCCTTATACCGTGGAGAACTAAATGAACATTTGGGACTCC  
TGGGGCCATATATAAGAGCAGAAGTTGAAGATAATATCATGTGTAACCTTCAGAAATCAGGCCTCTCGTCCCTATTCTTCT  
TATTCTAGCCTTATTTCTTATGAGGAAGATCAGAGGCAAGGAGCAGAACCTAGAAAAAACTTTGTCAAGCCTAATGAAAC  
CAAACTTACTTTTGGAAAGTGCAACATCATATGGCACCCACTAAAGATGAGTTTGAAGTCAAAGCCTGGGCTTATTTCT  
CTGATGTTGACCTGGAAAAAGATGTGCACTCAGGCCTGATTGGACCCCTTCTGGTCTGCCACACTAACACTGAACCTT  
GCTCATGGGAGACAAGTGACAGTACAGGAATTTGCTCTGTTTTTCACCATCTTTGATGAGACCAAAAGCTGGTACTTCAC  
TGAAAATATGGAAGAAACTGCAGGGCTCCCTGCAATATCCAGATGGAAGATCCCACTTTTAAAGAGAATTATCGCTTCC  
ATGCAATCAATGGCTACATAATGGATACACTACCTGGCTTAGTAATGGCTCAGGATCAAAGGATTCGATGGTATCTGCTC  
AGCATGGGCAGCAATGAAAACATCCATTCTATTCTTCACTGACATGTGTTCACTGTACGAAAAAAGAGGAGTATAA  
AATGGCACTGTACAATCTCTATCCAGGTGTTTTTGAGACAGTGGAAATGTTACCATCCAAAGCTGGAATTTGGCGGGTGG  
AATGCCTTATTTGGCGAGCATCTACATGCTGGGATGAGCACACTTTTTCTGGTGTACAGCAATAAGTGTGAGACTCCCTG  
GGAATGGCTTCTGGACACATTAGAGATTTTCAGATTACAGCTTCAGGACAATATGGACAGTGGGCCCCAAAGCTGGCCAG  
ACTTCATTATTCCGGATCAATCAATGCCTGGAGCACAAGGAGCCCTTTTCTGGATCAAGGTGGATCTGTTGGACCAA  
TGATTATTACGGCATCAAGACCAGGGTGGCCGTCAGAAGTTCTCCAGCCTCTACATCTCTCAGTTTATCATCATGTAT  
AGTCTTGATGGGAAGTGGCAGACTTATCGAGGAAATTCAGTGAACCTTAATGGTCTTCTTTGGCAATGTGGATTCT  
ATCTGGGATAAAAACAATATTTTTAACCTTCCAATTATTGCTCGATACATCCGTTTGACCCAACCTCATTATAGCATTC

GCAGCACTCTTCGCATGGAGTTGATGGGCTGTGATTTAAATAGTTGCAGCATGCCATTGGGAATGGAGAGTAAAGCAATA  
TCAGATGCACAGATTACTGCTTCATCCTACTTTACCAATATGTTTGCCACCTGGTCTCCTTCAAAAGCTCGACTTCACCT  
CCAAGGGAGGAGTAATGCTGGAGACCTCAGGTGAATAATCCAAAAGAGTGGCTGCAAGTGGACTTCCAGAAGACAATGA  
AAGTCACAGGAGTAACTACTCAGGGAGTAAATCTCTGCTTACCAGCATGTATGTGAAGGAGTTCCCTCATCTCCAGCAGT  
CAAGATGGCCATCAGTGGACTCTCTTTTTTCAGAATGGCAAAGTAAAGGTTTTTCAGGGAAATCAAGACTCCTTCACACC  
TGTGGTGAACCTCTCTAGACCCACCGTTACTGACTCGCTACCTTCGAATTCACCCCCAGAGTTGGGTGCACCAGATTGCCC  
TGAGGATGGAGTTCTGGGCTGCGAGGCACAGGACCTCTACTGACTCGAGAATAAAAGATCAGAGCTCTAGAGATCTGTG  
TGTGGTTTTTTGTGTGCGGCCGAGGAACCCCTAGTGATGGAGTTGGCCACTCCCTCTCTGCGCGCTCGCTCGCTCACT  
GAGGCCGGGCGCAAAAGGTCGCCCGACGCCCGGGCTTTGCCCGGGCGGCTCAGTGAGCGAGCGCGCGCAGCTGCCT  
GCAGGACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCTTGCTGGCGTTTTTCCATAGGCTCC  
GCCCCCTGACGAGCATCAGAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAGATACCAGGCG  
TTTTCCCTGGAGCTCCCTCGTGCGCTCTCTGTTCCGACCTGCCGCTTACCGGATACCTGTCCGCTTTCTCCCTTC  
GGGAAGCGTGGCGCTTTCTCATAGCTCAGCTGTAGGTATCTCAGTTCGGTGTAGGTGCTTCCGCTCCAAGCTGGGCTGTG  
TGCACGAACCCCCGTTTCAGCCCGACCGCTGCGCTTATCCGTAACCTATCGTCTTGAGTCCAACCCGGTAAGACACGAC  
TTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGATTCTTGAAGTG  
GTGGCCTAACTACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGAAAAAGAG  
TTGGTAGCTCTTGATCCGGCAAAACAAACCCGCTGGTAGCGGTGGTTTTTTTGTGTTGCAAGCAGCAGATTACGCGCAGA  
AAAAAGGATCTCAAGAAGATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGGAAACGAAACTCAGTTAAGGGAT  
TTTTGGTACATGATTCAAAAAAGGATCTTCACCTAGATCTCTTTTAAATTAATAAGTATTTAAATCAATCTAAAGTA  
TATATGAGTAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTCTGTTCA  
TCCATAGTTGCTGACTCCCCGTCGTGTAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGAT  
ACCGCGAGACCCACGCTCACCCTCCAGATTTATCAGCAATAAACAGCCAGCCGGAAGGGCCGAGCGCAGAAAGTGGTC  
CTGCAACTTTATCCGCTCCATCCAGTCTATTAATTTGTGCGGGAAGCTAGAGTAAGTAGTTCCGCCAGTTAATAGTTTG  
CGCAACGTTTGTGCTTGTGTCACAGCATCGTGTGTCACGCTCGTCTGTTGGTATGGCTTCATTACGCTCCGGTTCCCA  
ACGATCAAGGCGAGTTACATGATCCCCCATGTTGTGCAAAAAAGCGGTAGCTCCTTCGGTCCCTCCGATCGTTGTCAGAA  
GTAAGTTGGCCGAGTGTATCACTCATGTTTATGGCAGCACTGCATAATTTCTTACTGTCTATGCCATCCGTAAGATGC  
TTTTCTGTGACTGGTGTAGTACTCAACCAAGTCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCCCGCGTC  
AATACGGGATAATACCGCGCCATAGCAGAACTTTAAAAGTGCTCATCATTGGAACCGTTCTTCGGGGCGAAAACTCT  
CAAGGATCTTACCGCTGTGAGATCCAGTTGATTAACCCACTCGTGACCCCACTGATCTTCAGCATCTTTTACTTTTC  
ACCAGCGTTTCTGGGTGAGCAAAAAAGGAAGGCAAAAGCCGCGCAAAAAAGGGAATAAGGGCGACACGGAATCTTGAAT  
ACTCATACTCTTCTTTTTCAATATTATTGAAGCATTATCAGGGTTATTGTCTCATGAGCGGATACATATTGAATGTA  
TTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCACCTGACGCTTAAGAAACCATATTATC  
ATGACATTAACCTATAAAAAATAGGCGTATCAGGAGGCCCTTTTCGTCTCGCGCGTTTCGGTGATGACGGTGAAAACCTCTG  
ACACATGCAGCTCCCGGAGACGGTCACAGCTTGTCTGTAAGCGGATGCCGGGAGCAGACAAGCCCGTCAGGGCGCGTCAG  
CGGGTGTGGCGGGTGTGCGGGCTGGCTTAACCTATGCGGCATCAGAGCAGATTGTACTGAGAGTGACCATAAATGTGA  
AACGTTAATATTTTGTAAAAATTCGCGTTAAATTTTGTAAATCAGCTCATTTTTTTAACCAATAGGCCGAAATCGGCAA  
AATCCCTTATAAATCAAAAGATAGCCCCGAGATAGGGTGTAGTGTGTTCAGTTTGGAAACAGAGTCCACTATTAAAGA  
ACGTGGACTCCAACGTCAAAGGGCGAAAAACCGTCTATCAGGGCGATGGCCCACTACGTGAACCATCACCCAAATCAAGT  
TTTTTGGGGTCGAGGTGCCGTAAAGCACTAAATCGGAACCCCTAAAGGGAGCCCCGATTAGAGCTTGACGGGGAAAGCC  
GGCGAACGTGGCGAGAAAGGAAGGGAAGAAAGCGAAAGGAGCGGGCGCTAGGGCGCTGGCAAGTGATGCGGTCAAGCTGC  
GCGTAACCAACACACCCGCGCGCTTAATGCGCGCTACAGGGCGCGTACTATGGTTGCTTTGACGTATGCGGTGTGAAA  
TACCGCACAGATGCGTAAGGAGAAAAATACCGCATCAGGCCGTAAACCTGTCGGATCACCGGAAAGGACCCGTAAAGTGATA  
ATGATTATCATCTACATATCACAACGTGCGTGAGGCCATCAAACCAGTCAAATAATCAATTATGACGCAGGTATCGTA  
TTAATTGATCTGCATCAACTTAACGTAAAAACAACCTTCAGACAAATACAAATCAGCGACACTGAATACGGGGCAACCTCAT  
GTCAACGAAGAACAGAACCCGAGAACAAACCCGCAACATCCGCTTTCTTAACCAATGATTGAACAAATTAACATCG  
CTCTTGAGCAAAAAAGGTCCGGGAATTTCTCAGCTGGGTGATTGAAGCCTGCCGTCCGAGACTAACGTCAGAAAAGAGA  
GCATATACATCAATTAAGTGTATGAAGAATGAACATCCCGCTTCTTCCCTCCGAACAGGACGATATTGTAAATTCAT  
TAATTACGAGGGCATTGCAATTAAGTGTGAGTTTACCCTTTCTGACAGTGACAGACTGCGGTGTGGCTCTGTCA  
CAGACTAAATAGTTTGAATGATTAGCAGTTATGGTGTATCAGTCAACCACCGGGAATAATCCTTCATATTATTATCGTGC  
TTCACCAACGCTGCGCTCAATTGCTCTGAATGCTTCCAGAGACACCTTATGTTCTATACATGCAATTACAACATCAGGGTA  
ACTCATAGAAATGGTGTATTAAGCATATTTTTTACACGAATCAGATCCACGGAGGGATCATCAGCAGATTGTTCTTTAT  
TCATTTTGTGCTCCATGCGCTTGTCTCTCATCTAGCGGTAAAAATATTACTTCAAATCTTTCTGTATGAAGATTGTAGC  
ACGTTGGCTTACATACATCTGTGCGTTGTATTTCCCTCCAGAAATGCCAGCAGGACCGCACTTTGTTACGCAACCAATAC  
TATTAAGTGAACCATTCCTAATATTTGACATAAATCATCAACAAAACACAAGGAGGTGACAGCAGATTGAAACGATAAA  
AACGATAATGCAAACTACGCGCCCTCGTATCAGATGGAAGGTTTTTACCAATGGCTCAGGTTGCCATTTTTTAAAGAAATAT  
TCGATCAAGTGCGAAAGATTAGACTGTGAATGTTTATTCTGAACTAAAACGTCAACAGTCTCACATTATATTTAC  
TATCTAGCCACAGATAATATTCACATCGTGTAGAAAAAGGATAACACCGTGTAAATAAAGGACTTAAAAAGGTTGTAAA

TGTTAAATTCTCAAGAAACACGCATCTTATAGAAACGTCCTATGATAGGTTGAAATCAAGAGAAATCACATTTTCAGCAAT  
ACAGGGGAAAATCTTGCTAAAGCAGGAGTTTTCCGATGGGTTACAAATATCCATGAACATAAAAGATATTACTATACCTTT  
GATAATTCATTACTATTTACTGAGAGCATTTCAGAACTACACAAATCTTCCACGCTAAATCATAACGTCCGGTTTCTT  
CCGTGTCAGCACCGGGCGGTTGGCATAATGCAATACGTGTACGCGCTAAACCTGTGTGCATCGTTTTAATTATTCCTGG  
ACACTCCCGCAGAGAAGTTCCCGCTCAGGGCTGTGGACATAGTTAATCCGGGAATACAATGACGATTTCATCGCACCTGAC  
ATACATTAATAAATATTAACAATATGAAATTTCAACTCATTGTTTAGGGTTTGTTTAATTTTCTACACATACGATTCTGC  
GAACTTCAAAAAGCATCGGGAATAACACCATGAAAAAATGCTACTCGCTACTGCGCTGGCCCTGCTTATTACAGGATGT  
GCTCAACAGACGTTTACTGTTCAAAACAAACCGGCAGCAGTAGCACCAGGAAACCATCACCCTCATTTTCTTCGTTTC  
TGGAATTGGGCAGAGAAACTGTCGATGCAGCCAAAATTTGTGGCGGCGCAGAAAATGTTGTTAAACAGAAACCCAGC  
AAACATTTCGTAATGGATTGCTCGGTTTTATTACTTTAGGCATTTATACTCCGCTGGAAGCGGTGTGTATTGCTCACAA  
TAATTGCATGAGTTGCCCATCGCGATATGGGCAACTCTATCTGCACTGCTCATTAAATATACTTCTGGGTTCTTCCAGTT  
GTTTTTGCATAGTGATCAGCCTCTCTCTGAGGGTGAAATAATCCCGTTCAGCGGTGTCTGCCAGTCGGGGGGAGGCTGCA  
TTATCCACGCCCGAGGGCGGTGGTGGCTTCACGCACTGACTGACAGACTGCTTTGATGTGCAACCGACGACGACCGCGGC  
AACATCATCACGCAGAGCATCATTTTCAGCTTTAGCATCAGCTAACTCCTTCGTGTATTTGTCATCGAGCGCAGCAACAT  
CACGCTGACGCATCTGCATGTCAGTAAATGCCGCTTCGCCAGCTTCAGTTCTCTGGCATTTTGTGCTGGGCTTTG  
TAGGTAATGGCGTTATCACGGTAATGATTAACAGCCCATGACAGGCAGACGATGATGCAGATAACCAGAGCGGAGATAAT  
CGCGGTGACTCTGCTCATACATCAATCTCTGACCGTTCCGCCCGCTTCTTTGAATTTTGCAATCAGGCTGTACGCCCT  
ATGCTCGAACTGACCATAACCAGCGCCCGGCAGTGAAGCCAGATATTGCTGCAACGGTGCATTGCTGACGGATATCAC  
CAGCATCAATCATAGGTAAAGCGCCACGCTCCTTAATCTGCTGCAATGCCACAGCGTCTCTGACTTTTCGGAGAGAAGTCT  
TTCAGGCCAAGCTGCTTGCGGTAGGCATCCCAACCGGAAAGAAAGCTGGTAGCGTCCGGCGCCTGTTGATTGAGTTT  
TGGGTTTAGCGTGACAAGTTTGCGAGGGTGATCGGAGTAATCAGTAAATAGCTCTCCGCCCTACAATGACGTCATAACCAT  
GATTTCTGGTTTTCTGACGTCCGTTATCAGTTCCCTCCGACCACGCCAGCATATCGAGGAACGCCTTACGTTGATTATTG  
ATTTCTACCATCTTCTACTCCGGCTTTTTTAGCAGCGAAGCGTTTGATAAGCGAACCATCGAGTCAGTACCGATGTAGC  
CGATAAACACGCTCGTTATATAAGCGAGATTGCTACTTAGTCCGGCGAAGTCGAGAAGGTCAGAAATGAACAGGCGATA  
ATGGCGCATCGTTGCGTGCATTACTGTTTTGTAAACGCACCGCCATTATATCTGCCGGAAGGTACGCCATTGCAAA  
CGCAAGGATTGCCCGATGCTTGTCTTTGCGCGGAGAATGGCGGCCAACAGGTCATGTTTTCTGGCATCTTCATGT  
CTTACCCCCAATAAGGGGATTTGCTCTATTTAATTAGGAATAAGGTGCGATTACTGATAGAACAAATCCAGGCTACTGTGT  
TTAGTAATCAGATTTGTTGCTGACCGATATGCACGGGCAAAACGGCAGGAGGTGTTAGCGCGACCTCCTGCCACCCGCT  
TTCACGAAGGTCATGTGTAAAGGCGCAGCGTAATTAATTAATGAATTCAGGACAGACAGTGGCTACGGCTCAGTTT  
GGGTTGTGCTGTGCTGGGCGGCGATGACGCTGTACGCATTTGGTGATCCGGTTCTGCTTCCGGTATTGCTTAATTCA  
GCACAACGGAAAGAGCACTGGCTAACAGGCTCGCGACTCTTACGATTATCGACTCAATGCTCTTACCTGTTGTGCAG  
ATATAAAAAATCCCGAAACCGTTATGCAGGCTCTAACTATTACCTGCGAACTGTTTCGGGATTGCATTTTGACAGCTCT  
CTGCCTGCGATGGTTGGAGTTCCAGACGATACGTCGAAGTGACCAACTAGGCGGAATCGGTAGTAAGCGCCGCTCTTTT  
CATCTCACTACCAACAACGAGCGAATTAACCCATCGTTGAGTCAAATTTACCCAATTTTATTCAATAAGTCAATATCATGC  
CGTTAATATGTTGCCATCCGTGGCAATCATGCTGCTAACGTGTGACCGCATTCAAATGTTGTCTGCGATTGACTCTTCT  
TTGTGGCATTGCACCACCAGAGCGTCATACAGCGCTTAACAGTGCGTGACCGAGGTGGGTGGGTAAGGTTTGGGATTAG  
CATCGTCACAGCGCGATATGCTGCGCTGCTGGCATCCTTGAATAGCCGACGCCTTTCATCTTCCGCACTCTTCTCGA  
CAACTCTCCCCACAGCTCTGTTTTGGCAATATCAACCGCACGGCCTGTACCATGGCAATCTCTGCATCTTCCCCCGGC  
GTCGCGCACTACGGCAATAATCCGCATAAGCGAATGTTGCGAGCACTTGCAGTACCTTTGCCTTAGTATTTCTTCAAG  
CTGCCCCCTGCAGG

FIGURE 6

CGCCCCTGCAGGCAGCTGCGCGCTCGCTCGCTCACTGAGGCCGCCGGGCAA  
AGCCCGGGCGTCGGGCGACCTTTGGTCGCCCCGGCCTCAGTGAGCGAGCGAGC  
GCGCAGAGAGGGAGTGGCCAACTCCATCACTAGGGGTTCTTGCGGCCGCACG  
CGTGGTGGCGCGGGGTAAACTGGGAAAGTGATGTCGTGTAAGTGGCTCCGCCT  
TTTTCCCGAGGGTGGGGGAGAACCGTATATAAGTGCAGTAGTCCCGTGAAC  
GTTCTTTTTCGCAACGGGTTTGCCGCCCCGCGGCAGGTAAGTGCCAGGGAAT  
GTTTGTCTTAAATACCATCGCTCCAGGGAATGTTTGTCTTAAATACCATCT  
ACTGACACTGACATCCACTTTTTCTTTTCTCCACAGGTATCGATCCACCATGC  
AAATAGAGCTCTCCACCTGCTTCTTTCTGTGCCTTTTGCGATTCTGCTTTAGTG  
CCACCAGAAGATACTACCTGGGTGCAGTGGAAGTGTGATGGGACTATATGCA  
AAGTGATCTCGGTGAGCTGCCTGTGGACGCAAGATTTCTCCTAGAGTGCCA  
AAATCTTTTCCATTCAACACCTCAGTCGTGTACAAAAAGACTCTGTTTGTAGA  
ATTCACGGATCACCTTTTCAACATCGCTAAGCCAAGGCCACCCTGGATGGGTG  
TGCTAGGTCTACCATCCAGGCTGAGGTTTATGATACAGTGGTCATTACACTT  
AAGAACATGGCTTCCCATCCTGTGTCAGTCTTCATGCTGTTGGTGTATCCTACTG  
GAAAGCTTCTGAGGGAGCTGAATATGATGATCAGACCAGTCAAAGGGAGAA  
AGAAGATGATAAAGTCTTCCCTGGTGGAAAGCCATACATATGTCTGGCAGGTC  
CTGAAAGAGAAATGGTCCAATGGCCTCTGACCCACTGTGCCTTACCTACTCATA  
TCTTTCTCATGTGGACCTGGTAAAAGACTTGAATTCAGGCCTCATTGGAGCCC  
TACTAGTATGTAGAGAAGGGAGTCTGGCCAAGGAAAAGACACAGACCTTGCA  
CAAATTTATACTACTTTTGTCTGTATTTGATGAAGGGAAAAGTTGGCACTCAG  
AAACAAAGAACTCCTTGATGCAGGATAGGGATGCTGCATCTGCTCGGGCCTG  
GCCTAAAATGCACACAGTCAATGGTTATGTAAACAGGTCTCTGCCAGGTCTG  
ATTGGATGCCACAGGAAATCAGTCTATTGGCATGTGATTGGAATGGGCACCA  
CTCCTGAAGTGCACCTCAATATTCCTCGAAGGTACACATTTCTTGTGAGGAAC  
CATCGCCAGGCGTCTTGAAATCTCGCCAATAACTTTCTTACTGCTCAAAC  
ACTCTTGATGGACCTTGGACAGTTTCTACTGTTTTGTGATATCTCTTCCCACCA  
ACATGATGGCATGGAAGCTTATGTCAAAGTAGACAGCTGTCCAGAGGAACCC  
CAACTACGAATGAAAAATAATGAAGAAGCGGAAGACTATGATGATGATCTTA  
CTGATTCTGAAATGGATGTGGTCAGGTTTGATGATGACAACTCTCCTTCTTT  
ATCCAAATTCGCTCAGTTGCCAAGAAGCATCCTAAAACCTTGGGTACATTACAT  
TGCTGCTGAAGAGGAGGACTGGGACTATGCTCCCTTAGTCCTCGCCCCCGATG  
ACAGAAGTTATAAAAGTCAATATTTGAACAATGGCCCTCAGCGGATTGGTAG  
GAAGTACAAAAAAGTCCGATTTATGGCATAACACAGATGAAACCTTTAAGACT  
CGTGAAGCTATTCAGCATGAATCAGGAATCTTGGGACCTTTACTTTATGGGG  
AAGTTGGAGACACACTGTTGATTATATTTAAGAATCAAGCAAGCAGACCATA  
TAACATCTACCCTCACGGAATCACTGATGTCCGTCCTTTGTATTCAAGGAGAT  
TACCAAAAGGTGTAAACATTTGAAGGATTTTCCAATTCTGCCAGGAGAAAT  
ATTCAAATATAAATGGACAGTGACTGTAGAAGATGGGCCAACTAAATCAGAT  
CCTCGGTGCCTGACCCGCTATTACTCTAGTTTCGTTAATATGGAGAGAGATCT  
AGCTTCAGGACTCATTGGCCCTCTCCTCATCTGCTACAAAGAATCTGTAGATC  
AAAGAGGAAACCAGATAATGTCAGACAAGAGGAATGTCATCCTGTTTTCTGT  
ATTTGATGAGAACCGAAGCTGGTACCTCACAGAGAATATACAACGCTTTCTC

CCCA.TCCAGCTGGAGTGCAGCTTGAGGATCCAGAGTTCCAAGCCTCCAACA  
TCATGCACAGCATCAATGGCTATGTTTTTGATAGTTTGCAGTTGTCAGTTTGT  
TTGCATGAGGTGGCATACTGGTACATTCTAAGCATTGGAGCACAGACTGACT  
TCCTTTCTGTCTTCTTCTCTGGATATACCTTCAAACACAAAATGGTCTATGAA  
GACACACTCACCTATTCCCATTCTCAGGAGAACTGTCTTCATGTGATGGA  
AAACCCAGGTCTATGGATTCTGGGGTGCCACAACTCAGACTTTCGGAACAGA  
GGCATGACCGCCTTACTGAAGGTTTCTAGTTGTGACAAGAACAAGTGGTGATT  
ATTACGAGGACAGTTATGAAGATATTTTCAGCATACTTGCTGAGTAAAAACAA  
TGCCATTGAACCAAGAAGCTTCTCCCAGAATCCACCAGTCTTGAACGCCATC  
AACGCGAAATAACTCGTACTACTCTTCAGTCAGATCAAGAGGAAATTGACTA  
TGATGATACCATATCAGTTGAAATGAAGAAGGAAGATTTTGACATTTATGAT  
GAGGATGAAAATCAGAGCCCCCGCAGCTTTCAAAAGAAAACACGACACTATT  
TTATTGCTGCAGTGGAGAGGCTCTGGGATTATGGGATGAGTAGCTCCCCACA  
TGTTCTAAGAAACAGGGCTCAGAGTGGCAGTGTCCCTCAGTTCAAGAAAGTT  
GTTTTCCAGGAATTTACTGATGGCTCCTTTACTCAGCCCTTATACCGTGGAGA  
ACTAAATGAACATTTGGGACTCCTGGGGCCATATATAAGAGCAGAAGTTGAA  
GATAATATCATGGTAACTTTCAGAAATCAGGCCTCTCGTCCCTATTCTTCTA  
TTCTAGCCTTATTTCTTATGAGGAAGATCAGAGGCAAGGAGCAGAACCTAGA  
AAAAACTTTGTCAAGCCTAATGAAACCAAACTTACTTTTGAAAAGTGCAAC  
ATCATATGGCACCCACTAAAGATGAGTTTGACTGCAAAGCCTGGGCTTATTTCT  
TCTGATGTTGACCTGGAAAAAGATGTGCACTCAGGCCTGATTGGACCCCTTCT  
GGTCTGCCACACTAACACACTGAACCCTGCTCATGGGAGACAAGTGACAGTA  
CAGGAATTTGCTCTGTTTTTACCATCTTTGATGAGACCAAAAAGCTGGTACTT  
CACTGAAAATATGGAAAGAACTGCAGGGCTCCCTGCAATATCCAGATGGAA  
GATCCCACTTTTAAAGAGAATTATCGCTTCCATGCAATCAATGGCTACATAAT  
GGATACACTACCTGGCTTAGTAATGGCTCAGGATCAAAGGATTTCGATGGTAT  
CTGCTCAGCATGGGCAGCAATGAAAACATCCATTCTATTTCATTTCAGTGGACA  
TGTGTTCACTGTACGAAAAAAGAGGAGTATAAAATGGCAGACTGTACAATCTC  
TATCCAGGTGTTTTTGAGACAGTGGAAATGTTACCATCCAAGCTGGAATTTG  
GCGGGTGGAAATGCCTTATTGGCGAGCATCTACATGCTGGGATGAGCACACTT  
TTTCTGGTGTACAGCAATAAGTGTGAGACTCCCTGGGAATGGCTTCTGGACA  
CATTAGAGATTTTCAGATTACAGCTTCAGGACAATATGGACAGTGGGCCCCA  
AAGCTGGCCAGACTTCATTATTCCGGATCAATCAATGCCTGGAGCACCAAGG  
AGCCCTTTTCTTGGATCAAGGTGGATCTGTTGGCACCAATGATTATTCACGGC  
ATCAAGACCCAGGGTGCCCGTCAGAAGTTCTCCAGCCTCTACATCTCTCAGTT  
TATCATCATGTATAGTCTTGATGGGAAGAAGTGGCAGACTTATCGAGGAAAT  
TCCACTGGAACCTTAATGGTCTTCTTTGGCAATGTGGATTTCATCTGGGATAAA  
ACACAATATTTTTAACCCTCCAATTATTGCTCGATACATCCGTTTGCACCCAA  
CTCATTATAGCATTTCGACGACTCTTCGCATGGAGTTGATGGGCTGTGATTTA  
AATAGTTGCAGCATGCCATTGGGAATGGAGAGTAAAGCAATATCAGATGCAC  
AGATTACTGCTTCATCTTACTTTACCAATATGTTTGGCACCTGGTCTCCTTCAA  
AAGCTCGACTTCACCTCCAAGGGAGGAGTAATGCCTGGAGACCTCAGGTGAA  
TAATCCAAAAGAGTGGCTGCAAGTGGACTTCCAGAAGACAATGAAAGTCACA  
GGAGTAACTACTCAGGGAGTAAATCTCTGCTTACCAGCATGTATGTGAAGG  
AGTTCCTCATCTCCAGCAGTCAAGATGGCCATCAGTGGACTCTCTTTTTTCAG  
AATGGCAAAGTAAAGGTTTTTCAGGGAAATCAAGACTCCTTCACACCTGTGG

TGAACTCTCTAGACCCACCGTTACTGACTCGCTACCTTCGAATTCACCCCCAG  
AGTTGGGTGCACCAGATTGCCCTGAGGATGGAGGTTCTGGGCTGCGAGGCAC  
AGGACCTCTACTGACTCGAGCCTAATAAAGGAAATTTATTTTCATTGCAATAG  
TGTGTTGGTTTTTTGTGTGCGGCCGCAGGAACCCCTAGTGATGGAGTTGGCCA  
CTCCCTCTCTGCGCGCTCGCTCGCTCACTGAGGCCGGGCGACCAAAGGTCGCC  
CGACGCCCCGGGCTTTGCCCGGGCGGCCTCAGTGAGCGAGCGAGCGCGCAGCT  
GCCTGCAGGACAT

FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10





FIGURE 11



## SEQUENCE LISTING

<110> Couto, Linda B.  
Colosi, Peter C.

5

<120> ADENO-ASSOCIATED VECTORS FOR EXPRESSION OF FACTOR VIII  
BY TARGET  
CELLS

10 &lt;130&gt; AVIGEN-03743

<140> not yet assigned  
<141> 1999-07-30

15 <150> 60/125,974  
<151> 1999-03-24

<150> 60/104,994  
<151> 1998-10-20

20

&lt;160&gt; 14

&lt;170&gt; PatentIn Ver. 2.0

25 <210> 1  
<211> 59  
<212> DNA  
<213> Artificial Sequence30 <220>  
<223> Description of Artificial Sequence: Synthetic

<400> 1  
cccaagcttg cggccgcccc ggtgcccgcc ctaggcaggt aagtgccgtg tgtggttcc 59

35

<210> 2

<211> 59

<212> DNA

<213> Artificial Sequence

5

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 2

10 ccgctcgagc agagctctat ttgcatgggtg gaatcgatgc cgcggaacc acacacggc 59

<210> 3

<211> 103

<212> DNA

15 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

20 <400> 3

cccaagcttg cggccgccc ggtgccgcc ctaggcaggt aagtgccgtg tgtgggtccc 60  
gcggcatcga ttccaccatg caaatagagc tctgctcgag cgg 103

<210> 4

25 <211> 57

<212> DNA

<213> Artificial Sequence

<220>

30 <223> Description of Artificial Sequence: Synthetic

<400> 4

ttccgcggg cctggcctct ttacgggta tggcccttgc gtgccttgaa ttactga 57

35 <210> 5

<211> 57

<212> DNA

<213> Artificial Sequence

5 <220>

<223> Description of Artificial Sequence: Synthetic

<400> 5

10 gaatcgatac ctgtggagaa aaagaaaaag tggatgtcag tgtcagtaat tcaaggc 57

<210> 6

<211> 99

<212> DNA

<213> Artificial Sequence

15

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 6

20 tccccgcggg cctggcctct ttacgggtta tggcccttgc gtgccttgaa ttactgacac 60  
tgacatccac tttttctttt tctccacagg tatcgattc 99

<210> 7

<211> 100

25 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

30

<400> 7

aggggaatggtt tgttcttaaa taccatccag ggaatgtttg ttcttaaata ccatccaggg 60  
aatgtttggtt cttaaatacc atctacagtt attgggttaaa 100

35 <210> 8

<211> 59

<212> DNA

<213> Artificial Sequence

5 <220>

<223> Description of Artificial Sequence: Synthetic

<400> 8

ggaaagggtga tctgtgtgca gaaagactcg ctctaataata cttctttaac caataactg 59

10

<210> 9

<211> 144

<212> DNA

<213> Artificial Sequence

15

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 9

20 aggggaatggtt tggttcttaaa taccatccag ggaatgtttg ttcttaaata ccatccaggg 60  
aatggttggtt cttaaatacc atctacagtt attgggttaaa gaagtatatt agagcgagtc 120  
tttctgcaca cagatcacct ttcc 144

<210> 10

25 <211> 59

<212> DNA

<213> Artificial Sequence

<220>

30 <223> Description of Artificial Sequence: Synthetic

<400> 10

tgcgagaataa aagatcagag ctctagagat ctgtgtggtg gttttttgtg tgcggccgc 59

35 <210> 11

<211> 59

<212> DNA

<213> Artificial Sequence

5 <220>

<223> Description of Artificial Sequence: Synthetic

<400> 11

tcgagcggcc gcacacaaaa aaccaacaca cagatctcta gagctctgat cttttattc 59

10

<210> 12

<211> 63

<212> DNA

<213> Artificial Sequence

15

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 12

20 tcgagaataa aagatcagag ctctagagat ctgtgtgttg gttttttgtg tgcggccgct 60  
cga 63

<210> 13

<211> 11933

25 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: synthetic

30

<400> 13

cagctgcgcg ctgcctcgct cactgaggcc gcccgggcaa agcccgggcg tcgggcgacc 60  
tttggtcgcc cggcctcagt gagcgagcga gcgcgcagag agggagtggc caactccatc 120  
actaggggtt cctgcggccg ccaggggaat gtttgttctt aaataccatc cagggaatgt 180  
35 ttgttcttaa ataccatcta gggaatgttt gttcttaaata accatctaca gttattggtt 240

aaagaagtat attagagcga gtctttctgc acacagatca cctttccggg tgccgcccct 300  
 aggcaggtaa gtgccgtgtg tggttcccg gggcctggcc tctttacggg ttatggccct 360  
 tgcgtgcctt gaattactga cactgacatc cactttttct ttttctccac aggtatcgat 420  
 tccaccatgc aaatagagct ctccacctgc ttctttctgt gccttttgcg attctgcttt 480  
 5 agtgccacca gaagatacta cctgggtgca gtggaactgt catgggacta tatgcaaagt 540  
 gatctcggtg agctgcctgt ggacgcaaga tttcctccta gagtgccaaa atcttttcca 600  
 ttcaacacct cagtcgtgta caaaaagact ctgtttgtag aattcacgga tcaccttttc 660  
 aacatcgcta agccaaggcc accctggatg ggtctgctag gtcctaccat ccaggctgag 720  
 gtttatgata cagtgggtcat tacacttaag aacatggctt cccatcctgt cagtcttcat 780  
 10 gctgttggtg tctcctactg gaaagcttct gagggagctg aatatgatga tcagaccagt 840  
 caaagggaga aagaagatga taaagtcttc cctgggtggaa gccatacata tgtctggcag 900  
 gtcctgaaag agaatggtcc aatggcctct gacctactgt gccttaccta ctcatatctt 960  
 tctcatgtgg acctggtaaa agacttgaat tcaggcctca ttggagccct actagtatgt 1020  
 agagaaggga gtctggccaa ggaaaagaca cagaccttgc acaaatttat actacttttt 1080  
 15 gctgtatttg atgaaggga aagttggcac tcagaaacaa agaactcctt gatgcaggat 1140  
 agggatgctg catctgctcg ggctggcctt aaaatgcaca cagtcaatgg ttatgtaaac 1200  
 aggtctctgc caggtctgat tggatgccac aggaaatcag tctattggca tgtgattgga 1260  
 atgggcacca ctctgaagt gcactcaata ttctcgaag gtcacacatt tcttgtgagg 1320  
 aaccatcgcc aggcgtcctt ggaaatctcg ccaataactt tccttactgc tcaaactc 1380  
 20 ttgatggacc ttggacagtt tctactgttt tgtcatatct ctcccacca acatgatggc 1440  
 atggaagctt atgtcaaagt agacagctgt ccagaggaac cccaactacg aatgaaaaat 1500  
 aatgaagaag cggaagacta tgatgatgat cttactgatt ctgaaatgga tgtggtcagg 1560  
 tttgatgatg acaactctcc ttcttttacc caaattcgct cagttgccaa gaagcatcct 1620  
 aaaacttggg tacattacat tgctgctgaa gaggaggact gggactatgc tcccttagtc 1680  
 25 ctgcccccg atgacagaag ttataaaagt caatatttga acaatggccc tcagcggatt 1740  
 ggtaggaagt acaaaaaagt ccgatttatg gcatacacag atgaaacctt taagactcgt 1800  
 gaagctattc agcatgaatc aggaatcttg ggacctttac tttatgggga agttggagac 1860  
 aactgttga ttatatatta gaatcaagca agcagaccat ataacatcta ccctcacgga 1920  
 atcactgatg tccgtccttt gtattcaagg agattaccaa aaggtgtaaa acatttgaag 1980  
 30 gattttccaa ttctgccagg agaaatattc aaatataaat ggacagtga tgtagaagat 2040  
 gggccaacta aatcagatcc tcgggtgctg acccgctatt actctagttt cgtaaatatg 2100  
 gagagagatc tagcttcagg actcattggc cctctcctca tctgctacaa agaactctgta 2160  
 gatcaaagag gaaaccagat aatgtcagac aagaggaatg tcctcctgtt ttctgtattt 2220  
 gatgagaacc gaagctggta cctcacagag aatatacaac gctttctccc caatccagct 2280  
 35 ggagtgcagc ttgaggatcc agagttccaa gcctccaaca tcatgcacag catcaatggc 2340

tatgtttttg atagtttgca gttgtcagtt tgtttgcatg aggtggcata ctggtacatt 2400  
 ctaagcattg gagcacagac tgacttcctt tctgtcttct tctctggata taccttcaaa 2460  
 cacaaaatgg tctatgaaga cacactcacc ctattcccat tctcaggaga aactgtcttc 2520  
 atgtcgatgg aaaaccagg tctatggatt ctgggggtgcc acaactcaga ctttcggaac 2580  
 5 agaggcatga ccgccttact gaaggtttct agttgtgaca agaactctgg tgattattac 2640  
 gaggacagtt atgaagatat ttcagcatat ttgctgagta aaaacaatgc cattgaacca 2700  
 agaagcttcg aaataactcg tactactctt cagtcagatc aagaggaaat tgactatgat 2760  
 gataccatat cagttgaaat gaagaaggaa gattttgaca tttatgatga ggatgaaaat 2820  
 cagagcccc gcagctttca aaagaaaaca cgacactatt ttattgctgc agtggagagg 2880  
 10 ctctgggatt atgggatgag tagctcccca catgttctaa gaaacagggc tcagagtggc 2940  
 agtgtccctc agttcaagaa agttgttttc caggaattta ctgatggctc ctttactcag 3000  
 cccttatacc gtggagaact aaatgaacat ttgggactcc tggggccata tataagagca 3060  
 gaagttgaag ataatatcat ggtaactttc agaaatcagg cctctcgtcc ctattccttc 3120  
 tattctagcc ttatttctta tgaggaagat cagaggcaag gagcagaacc tagaaaaaac 3180  
 15 tttgtcaagc ctaatgaaac caaaacttac ttttggaag tgcaacatca tatggcacc 3240  
 actaaagatg agtttgactg caaagcctgg gcttatttct ctgatgttga cctggaaaaa 3300  
 gatgtgcact caggcctgat tggaccctt ctggtctgcc acactaacac actgaaccct 3360  
 gctcatggga gacaagtgc agtacaggaa ttgctctgt ttttcacat ctttgatgag 3420  
 accaaaagct ggtacttcac tgaaaatat gaaagaaact gcagggtcc ctgcaatatc 3480  
 20 cagatggaag atcccacttt taaagagaat tatcgttcc atgcaatcaa tggtacata 3540  
 atggatacac tacctggctt agtaatggct caggatcaaa ggattcgatg gtatctgctc 3600  
 agcatgggca gcaatgaaaa catccattct attcatttca gtggacatgt gttcactgta 3660  
 cgaaaaaaag aggagtataa aatggcactg tacaatctct atccaggtgt ttttgagaca 3720  
 gtggaaatgt taccatccaa agctggaatt tggcgggtgg aatgccttat tggcgagcat 3780  
 25 ctacatgctg ggatgagcac actttttctg gtgtacagca ataagtgtca gactcccctg 3840  
 ggaatggctt ctggacacat tagagatttt cagattacag cttcaggaca atatggacag 3900  
 tgggccccaa agctggccag acttcattat tccggatcaa tcaatgcctg gagcaccaag 3960  
 gagccctttt cttggatcaa ggtggatctg ttggcaccaa tgattattca cggcatcaag 4020  
 acccaggggtg ccgctcagaa gttctccagc ctctacatct ctcagtttat catcatgtat 4080  
 30 agtcttgatg ggaagaagtg gcagacttat cgaggaaatt cactggaac cttaatggct 4140  
 ttctttggca atgtggatct atctgggata aaacacaata tttttaacc tccaattatt 4200  
 gctcgatata tccgtttgca cccaactcat tatagcatc gcagcactct tcgcatggag 4260  
 ttgatgggct gtgatttaaa tagttgcagc atgccattgg gaatggagag taaagcaata 4320  
 tcagatgcac agattactgc ttcacacctac ttaccaata tgtttgccac ctggtctcct 4380  
 35 tcaaaagctc gacttcacct ccaagggagg agtaatgcct ggagacctca ggtgaataat 4440

```

ccaaaagagt ggctgcaagt ggacttccag aagacaatga aagtcacagg agtaactact 4500
cagggagtaa aatctctgct taccagcatg tatgtgaagg agttcctcat ctccagcagt 4560
caagatggcc atcagtggac tctctttttt cagaatggca aagtaaagggt ttttcagggga 4620
aatcaagact ctttcacacc tgtggtgaac tctctagacc caccgttact gactcgctac 4680
5  cttcgaattc acccccagag ttgggtgcac cagattgccc tgaggatgga ggttctgggc 4740
tgcgaggcac aggacctcta ctgactcgag aataaaagat cagagctcta gagatctgtg 4800
tgttggtttt ttgtgtgcgg ccgcaggaac ccctagtgat ggagttggcc actccctctc 4860
tgcgcgctcg ctcgctcact gaggccgggc gaccaaagggt cgcgcgacgc ccgggctttg 4920
cccgggcggc ctgagtgcgc gagcgagcgc gcagctgcct gcaggacatg tgagcaaaaag 4980
10 gccagcaaaa ggccaggaac cgtaaaaagg ccgcgttgct ggcgtttttc cataggctcc 5040
gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga aaccgcacag 5100
gactataaag ataccaggcg tttccccctg gaagctccct cgtgcgctct cctgttccga 5160
ccctgccgct taccggatac ctgtccgcct ttctcccttc gggaagcgtg gcgctttctc 5220
atagctcacg ctgtaggtat ctgagttcgg tgtaggtcgt tcgctccaag ctgggctgtg 5280
15 tgcacgaacc ccccgttcag ccgcaccgct gcgccttata cggtaactat cgtcttgagt 5340
ccaacccggt aagacacgac ttatcgccac tggcagcagc cactggtaac aggattagca 5400
gagcgaggta tgtaggcggg gctacagagt tcttgaagtg gtggcctaac tacggctaca 5460
ctagaaggac agtatttggg atctgcgctc tgctgaagcc agttaccttc ggaaaaagag 5520
ttggtagctc ttgatccggc aaacaaacca ccgctggtag cgggtggtttt tttgtttgca 5580
20 agcagcagat tacgcgcaga aaaaaaggat ctcaagaaga tcctttgatc ttttctacgg 5640
ggctctgacgc tcagtggaac gaaaactcac gttaagggat tttggtcatt agattatcaa 5700
aaaggatctt cacctagatc ctttttaatt aaaaatgaag ttttaaatca atctaaagta 5760
tatatgagta aacttgggtc gacagttacc aatgcttaat cagtgaggca cctatctcag 5820
cgatctgtct atttcgttca tccatagttg cctgactccc cgtcgtgtag ataactacga 5880
25 tacgggaggg cttaccatct ggccccagtg ctgcaatgat accgcgagac ccacgctcac 5940
cggctccaga tttatcagca ataaaccagc cagccggaag ggccgagcgc agaagtgggc 6000
ctgcaacttt atccgcctcc atccagtcta ttaattgttg ccgggaagct agagtaagta 6060
gttcgccagt taatagtttg cgcaacgttg ttgccattgc tacaggcatc gtggtgtcac 6120
gctcgtcgtt tggatgggt tcattcagct ccggttccca acgatcaagg cgagttacat 6180
30 gatcccccatt gttgtgcaaa aaagcgggta gctccttcgg tcctccgata gttgtcagaa 6240
gtaagttggc cgcagtgtta tcaactcatg ttatggcagc actgcataat tctcttactg 6300
tcatgccatc cgtaagatgc ttttctgtga ctggtgagta ctcaaccaag tcattctgag 6360
aatagtgtat gcggcgaccg agttgctctt gccggcgctc aatacgggat aataccgcgc 6420
cacatagcag aactttaaaa gtgctcatca ttggaaaacg ttcttcgggg cgaaaactct 6480
35 caaggatctt accgctgttg agatccagtt cgatgtaacc cactcgtgca cccaactgat 6540

```

cttcagcatc ttttactttc accagcggtt ctgggtgagc aaaaacagga aggcaaaatg 6600  
 ccgcaaaaaa gggaataagg gcgacacgga aatgttgaat actcatactc ttcctttttc 6660  
 aatattattg aagcatttat cagggttatt gtctcatgag cggatacata tttgaatgta 6720  
 tttagaaaaa taaacaaata ggggttccgc gcacatttcc ccgaaaagtg ccacctgacg 6780  
 5 tctaagaaac cattattatc atgacattaa cctataaaaa taggcgtatc acgaggccct 6840  
 ttcgtctcgc gcgtttcggg gatgacggtg aaaacctctg acacatgcag ctcccggaga 6900  
 cggtcacagc ttgtctgtaa gcggatgccg ggagcagaca agcccgtcag ggcgcgtcag 6960  
 cgggtgttgg cgggtgtcgg ggctggctta actatgcggc atcagagcag attgtactga 7020  
 gagtgcacca taaaattgta aacgttaata ttttgtaaa attcgcgtta aatttttgtt 7080  
 10 aaatcagctc attttttaac caataggccg aaatcggcaa aatcccttat aaatcaaaag 7140  
 aatagcccgga gatagggttg agtggtgttc cagtttgga caagagtcca ctattaaaga 7200  
 acgtggactc caacgtcaaa gggcgaaaaa ccgtctatca gggcgatggc ccactacgtg 7260  
 aaccatcacc caaatcaagt tttttggggg cgagggtgccg taaagcacta aatcgggaacc 7320  
 ctaaagggag ccccgattt agagcttgac ggggaaagcc ggcgaaacgtg gcgagaaagg 7380  
 15 aaggaagaa agcgaaagga gcgggcgcta gggcgctggc aagtgtagcg gtcacgctgc 7440  
 gcgtaaccac cacacccgcc gcgcttaatg cgccgctaca gggcgctac tatggttgct 7500  
 ttgacgtatg cgggtgtgaaa taccgcacag atgcgtaagg agaaaatacc gcatcaggcc 7560  
 gtaacctgtc ggatcaccgg aaaggacccg taaagtata atgattatca tctacatatc 7620  
 acaacgtgcg tggaggccat caaacacgt caaataatca attatgacgc aggtatcgta 7680  
 20 ttaattgatc tgcataact taacgtaaaa acaacttcag acaatacaaa tcagcgacac 7740  
 tgaatacggg gcaacctcat gtcaacgaag aacagaacct gcagaacaac aaccgcgaac 7800  
 atccgctttc ctaaccaaact gattgaacaa attaacatcg ctcttgagca aaaaggggtc 7860  
 gggaatttct cagcctgggt cattgaagcc tgccgtcggg gactaacgtc agaaaagaga 7920  
 gcatatacat caattaaaag tgatgaagaa tgaacatccc gcgttcttcc ctccgaacag 7980  
 25 gacgatattg taaattcact taattacgag ggcattgcag taattgagtt gcagttttac 8040  
 cactttcctg acagtgcag actgcgtgtt ggctctgtca cagactaaat agtttgaatg 8100  
 attagcagtt atggtgatca gtcaaccacc aggaataat ccttcataatt attatcgtgc 8160  
 ttcaccaacg ctgcctcaat tgctctgaat gcttcagag acaccttatg ttctatacat 8220  
 gcaattacaa catcagggtg actcatagaa atggtgctat taagcatatt ttttacacga 8280  
 30 atcagatcca cggagggatc atcagcagat tggtctttat tcattttgtc gctccatgcg 8340  
 cttgctcttc atctagcggg taaaatatta cttcaaatct ttctgtatga agatttgagc 8400  
 acgttggcct tacatacatc tgtcgggtgt atttccctcc agaatgccag caggaccgca 8460  
 ctttgttacg caaccaatac tattaagtga aaacattcct aatatttgac ataaatcatc 8520  
 aacaaaacac aaggaggtca gaccagattg aaacgataaa aacgataatg caaactacgc 8580  
 35 gccctcgtat cacatggaag gttttaccaa tggtcaggt tgccattttt aaagaaatat 8640

tcgatcaagt gcgaaaagat ttagactgtg aattgtttta ttctgaacta aaacgtcaca 8700  
 acgtctcaca ttatatattac tatctagcca cagataatat tcacatcgtg ttagaaaacg 8760  
 ataacaccgt gttaataaaa ggacttaaaa aggttgtaaa tgttaaattc tcaagaaaca 8820  
 cgcattcttat agaaacgtcc tatgataggt tgaaatcaag agaaatcaca tttcagcaat 8880  
 5 acagggaaaa tcttgctaaa gcaggagttt tccgatgggt tacaaatata catgaacata 8940  
 aaagatatta ctataccttt gataattcat tactatttac tgagagcatt cagaacacta 9000  
 caciaaatctt tccacgctaa atcataacgt ccggtttctt ccgtgtcagc accggggcgt 9060  
 tggcataatg caatacgtgt acgcgctaaa ccctgtgtgc atcgttttaa ttattcccgg 9120  
 aactcccgc agagaagttc cccgtcaggg ctgtggacat agttaatccg ggaatacaat 9180  
 10 gacgattcat cgcacctgac atacattaat aaatattaac aatatgaaat ttcaactcat 9240  
 tgtttagggt ttgtttaatt ttctacacat acgattctgc gaacttcaaa aagcatcggg 9300  
 aataacacca tgaaaaaaat gctactcgt actgcgctgg ccctgcttat tacaggatgt 9360  
 gctcaacaga cgtttactgt tcaaaaacaaa ccggcagcag tagcaccaaa ggaaaccatc 9420  
 acccatcatt tcttcgtttc tggaattggg cagaagaaaa ctgtcgtatc agccaaaatt 9480  
 15 tgtggcggcg cagaaaaatgt tgtaaaaaca gaaaccagc aaacattcgt aaatggattg 9540  
 ctcggtttta ttactttagg catttatact ccgctggaag cgcgtgtgta ttgctcacia 9600  
 taattgcatg agttgcccac cgcgatatgg gcaactctat ctgcactgct cattaatata 9660  
 cttctgggtt ccttcaggtt gtttttgcac agtgatcagc ctctctctga gggtgaaata 9720  
 atcccgttca gcggtgtctg ccagtcgggg ggaggctgca ttatccacgc cggaggcggg 9780  
 20 ggtggcttca cgcactgact gacagactgc tttgatgtgc aaccgacgac gaccagcggc 9840  
 aacatcatca cgcagagcat cattttcagc tttagcatca gctaactcct tcgtgtatct 9900  
 tgcacgagc gcagcaacat cacgctgacg catctgcatg tcagtaattg ccgcgttcgc 9960  
 cagcttcagt tctctggcat ttttgcgcg ctgggctttg taggtaattg cgttatcacg 10020  
 gtaatgatta acagcccatg acaggcagac gatgatgcag ataaccagag cggagataat 10080  
 25 cgcggtgact ctgctcaca atcaatctct ctgaccgttc cgcgcgttc tttgaatttt 10140  
 gcaatcaggc tgcagcctt atgctcgaac tgaccataac cagcgcccg cagtgaagcc 10200  
 cagatattgc tgcaacggtc gattgcctga cggatatcac cacgatcaat cataggtaaa 10260  
 gcgccacgct ccttaatctg ctgcaatgcc acagcgtcct gacttttcgg agagaagtct 10320  
 ttcaggccaa gctgcttgcg gtaggcatcc caccaacggg aaagaagctg gtagcgtccg 10380  
 30 gcgcctgttg atttgagttt tgggttttagc gtgacaagtt tgcgagggtg atcggagtaa 10440  
 tcagtaata gctctccgcc tacaatgacg tcataaaccat gatttctggg tttctgacgt 10500  
 ccgttatcag ttccctccga ccacgccagc atatcgagga acgccttacg ttgattattg 10560  
 atttctacca tcttctactc cggctttttt agcagcgaag cgtttgataa gcgaaccaat 10620  
 cgagtcagta ccgatgtagc cgataaacac gctcgttata taagcgagat tgctacttag 10680  
 35 tccggcgaag tcgagaaggt cacgaatgaa ccaggcgata atggcgcaca tcgttgcgctc 10740

gattactgtt tttgtaaagc caccgccatt atatctgccg cgaaggtagc ccattgcaaa 10800  
 cgcaaggatt gccccgatgc cttgttcctt tgccgcgaga atggcggcca acaggatcatg 10860  
 tttttctggc atcttcatgt cttaccccca ataaggggat ttgctctatt taattaggaa 10920  
 taaggatgat tactgataga acaaattccag gctactgtgt ttagtaataca gattttgttcg 10980  
 5 tgaccgatat gcacgggcaa aacggcagga ggttgtagc gcgacctct gccacccgct 11040  
 ttcacgaagg tcatgtgtaa aaggccgcag cgtaactatt actaatgaat tcaggacaga 11100  
 cagtggctac ggctcagttt ggggttggtc gttgctgggc ggcatgacg cctgtacgca 11160  
 tttggtgatc cgggttctgt tccggtattc gcttaattca gcacaacgga aagagcactg 11220  
 gctaaccagg ctgcgcgact cttcacgatt atcgactcaa tgctcttacc tgttggtgcag 11280  
 10 atataaaaaa tcccgaacc gttatgcagg ctctaactat tacctgcgaa ctgtttcggg 11340  
 attgcatttt gcagacctct ctgcctgcga tgggtggagt tccagacgat acgtcgaagt 11400  
 gaccaactag gcggaatcgg tagtaagcgc cgcctctttt catctcacta ccacaacgag 11460  
 cgaattaacc catcggttag tcaaattttac ccaattttat tcaataagtc aatatcatgc 11520  
 cgtaaatatg ttgcatccg tggcaatcat gctgctaacg tgtgaccgca ttcaaatgt 11580  
 15 tgtctgcgat tgactcttct ttgtggcatt gcaccaccag agcgctatac agcggcttaa 11640  
 cagtgcgtga ccagggtggg tgggtaagg tgggattag catcgtcaca gcgcgatatg 11700  
 ctgcgcttgc tggcatcctt gaatagccga cgcctttgca tcttcgcac tctttctcga 11760  
 caactctccc ccacagctct gttttggcaa tatcaaccgc acggcctgta ccatggcaat 11820  
 ctctgcactc tgcgcccggc gtcgcggcac tacggcaata atccgcataa gcgaatgttg 11880  
 20 cgagcacttg cagtacctt gccttagtat ttccttcaag ctgcccctgc agg 11933

<210> 14

<211> 4999

<212> DNA

25 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

30 <400> 14

cgcccctgca ggcagctgcg cgctcgctcg ctactgagg ccgcccgggc aaagcccggg 60  
 cgtcgggcga cctttggctg cccggcctca gtgagcgagc gagcgcgag agagggagtg 120  
 gccaaactcca tcaatagggg ttccctgcggc cgcacgcgtg gtggcgcggg gtaaaactggg 180  
 aaagtgatgt cgtgtactgg ctccgccttt ttcccgaggg tgggggagaa ccgtatataa 240  
 35 gtgcagtagt cgccgtgaac gttctttctc gcaacgggtt tgccgccccg cggcaggtaa 300

gtgccagggga atgtttgttc ttaaataacca tcgctccagg gaatgtttgt tcttaaatac 360  
 catctactga cactgacatc cactttttct ttttctccac aggtatcgat ccaccatgca 420  
 aatagagctc tccacctgct tctttctgtg ccttttgcca ttctgcttta gtgccaccag 480  
 aagatactac ctgggtgcag tggaaactgtc atgggactat atgcaaagtg atctcggtga 540  
 5 gctgcctgtg gacgcaagat ttctctctag agtgccaaaa tcttttccat tcaacacctc 600  
 agtcgtgtac aaaaagactc tgtttgtaga attcacggat caccttttca acatcgctaa 660  
 gccaaggcca ccctggatgg gtctgctagg tcctaccatc caggctgagg tttatgatac 720  
 agtggtcatt acacttaaga acatggcttc ccactctgtc agtcttcatg ctgttggtgt 780  
 atcctactgg aaagcttctg agggagctga atatgatgat cagaccagtc aaagggagaa 840  
 10 agaagatgat aaagtcttcc ctggtggaag ccatacatat gtctggcagg tcctgaaaga 900  
 gaatggtcca atggcctctg acccactgtg ccttacctac tcatatcttt ctcatgtgga 960  
 cctggtaaaa gacttgaatt caggcctcat tggagcccta ctagtatgta gagaagggag 1020  
 tctggccaag gaaaagacac agaccttgca caaatattata ctactttttg ctgtatttga 1080  
 tgaagggaaa agttggcact cagaaacaaa gaactccttg atgcaggata gggatgctgc 1140  
 15 atctgctcgg gcctggccta aaatgcacac agtcaatggt tatgtaaaca ggtctctgcc 1200  
 aggtctgatt ggatgccaca ggaaatcagt ctattggcat gtgattggaa tgggcaccac 1260  
 tcctgaagtg cactcaatat tcctcgaagg tcacacattt cttgtgagga accatcgcca 1320  
 ggcgtccttg gaaatctcgc caataacttt ccttactgct caaacactct tgatggacct 1380  
 tggacagttt ctactgtttt gtcatatctc tccccaccaa catgatggca tgggaagctta 1440  
 20 tgtcaaagta gacagctgtc cagaggaacc ccaactacga atgaaaaata atgaagaagc 1500  
 ggaagactat gatgatgac ttactgattc tgaaatggat gtggtcagggt ttgatgatga 1560  
 caactctcct tcctttatcc aaattcgctc agttgccaaag aagcatccta aaacttgggt 1620  
 acattacatt gctgctgaag agggaggactg ggactatgct cccttagtcc tcgccccga 1680  
 tgacagaagt tataaaagtc aatatttgaa caatggcct cagcggattg gtaggaagta 1740  
 25 caaaaaagtc cgatttatgg catacacaga tgaaaccttt aagactcgtg aagctattca 1800  
 gcatgaatca ggaatcttgg gacctttact ttatggggaa gttggagaca cactgttgat 1860  
 tatatttaag aatcaagcaa gcagaccata taacatctac cctcacggaa tcatgatgt 1920  
 ccgtcctttg tattcaagga gattaccaa aggtgtaaaa catttggaagg attttccaat 1980  
 tctgccagga gaaatattca aatataaatg gacagtgact gtagaagatg ggccaactaa 2040  
 30 atcagatcct cgggtgctga cccgctatta ctctagtttc gttaatatgg agagagatct 2100  
 agcttcagga ctcatgggc ctctcctcat ctgctacaaa gaatctgtag atcaaagagg 2160  
 aaaccagata atgtcagaca agaggaatgt catcctgttt tctgtatttg atgagaaccg 2220  
 aagctggtac ctacagaga atatacaacg ctttctcccc aatccagctg gagtgcagct 2280  
 tgaggatcca gagttccaag cctccaacat catgcacagc atcaatggct atgtttttga 2340  
 35 tagtttcgag ttgtcagttt gtttgcagta ggtggcatatc tggtacatc taagcattgg 2400

agcacagact gacttccttt ctgtcttctt ctctggatat accttcaaac acaaaatggg 2460  
 ctatgaagac acactcaccc tattccatt ctcaggagaa actgtcttca tgtcgatgga 2520  
 aaacccaggt ctatggattc tggggtgcc aactcagac tttcggaaca gaggcattgac 2580  
 cgccttactg aaggtttcta gttgtgacaa gaacactggg gattattacg aggacagtta 2640  
 5 tgaagatatt tcagcatact tgctgagtaa aaacaatgcc attgaaccaa gaagcttctc 2700  
 ccagaatcca ccagtcttga aacgccatca acgcgaaata actcgtacta ctcttcagtc 2760  
 agatcaagag gaaattgact atgatgatac catatcagtt gaaatgaaga aggaagattt 2820  
 tgacatttat gatgaggatg aaaatcagag ccccccgcagc tttcaaaaga aaacacgaca 2880  
 ctatttttatt gctgcagtgg agaggctctg ggattatggg atgagtagct ccccacatgt 2940  
 10 tctaagaaac agggctcaga gtggcagtgt ccctcagttc aagaaagttg ttttccagga 3000  
 atttactgat ggctccttta ctcagccctt ataccgtgga gaactaaatg aacatttggg 3060  
 actcctgggg ccatatataa gagcagaagt tgaagataat atcatggtaa ctttcagaaa 3120  
 tcaggcctct cgtccctatt ccttctattc tagccttatt tcttatgagg aagatcagag 3180  
 gcaaggagca gaacctagaa aaaactttgt caagcctaata gaaacaaaaa cttacttttg 3240  
 15 gaaagtgcaa catcatatgg caccactaa agatgagttt gactgcaaag cctgggctta 3300  
 tttctctgat gttgacctgg aaaaagatgt gcactcaggc ctgattggac cccttctggg 3360  
 ctgccacact aacacactga accctgctca tgggagacaa gtgacagtac aggaatttgc 3420  
 tctgtttttc accatctttg atgagaccaa aagctggtac ttcactgaaa atatggaaag 3480  
 aaactgcagg gctccctgca atatccagat ggaagatccc acttttaaag agaattatcg 3540  
 20 cttccatgca atcaatggct acataatgga tacactacct ggcttagtaa tggctcagga 3600  
 tcaaaggatt cgatggatc tgctcagcat gggcagcaat gaaaacatcc attctattca 3660  
 tttcagtgga catgtgttca ctgtacgaaa aaaagaggag tataaaatgg cactgtacaa 3720  
 tctctatcca ggtgtttttg agacagtgga aatgttacca tccaaagctg gaatttggcg 3780  
 ggtggaatgc cttattggcg agcatctaca tgctgggatg agcacacttt ttctgggtgta 3840  
 25 cagcaataag tgctcagact ccctgggaat ggcttctgga cacattagag attttcagat 3900  
 tacagcttca ggacaatatg gacagtgggc cccaaagctg gccagacttc attattccgg 3960  
 atcaatcaat gcctggagca ccaaggagcc cttttcttgg atcaagggtg atctgttggc 4020  
 accaatgatt attcacggca tcaagaccca ggggtgcccgt cagaagttct ccagcctcta 4080  
 catctctcag tttatcatca tgtatagtct tgatgggaag aagtggcaga cttatcgagg 4140  
 30 aaattccact ggaaccttaa tgggtcttct tggcaatgtg gattcatctg ggataaaaca 4200  
 caatattttt aacctccaa ttattgctcg atacatccgt ttgcaccaa ctcatatag 4260  
 cattcgcagc actcttcgca tggagtgtgat gggctgtgat ttaaatagtt gcagcatgcc 4320  
 attgggaatg gagagtaaag caatatcaga tgcacagatt actgcttcat cctactttac 4380  
 caatatgttt gccacctggg ctcttcaaaa agctcgactt cacctccaag ggaggagtaa 4440  
 35 tgcctggaga cctcaggtga ataatccaaa agagtggctg caagtggact tccagaagac 4500

aatgaaagtc acaggagtaa ctactcaggg agtaaaatct ctgcttacca gcatgtatgt 4560  
gaaggagttc ctcattctcca gcagtcaaga tggccatcag tggactctct tttttcagaa 4620  
tggcaaagta aagggttttc agggaaatca agactccttc acacctgtgg tgaactctct 4680  
agacccaccg ttactgactc gctaccttcg aattcacccc cagagttggg tgcaccagat 4740  
5 tgccctgagg atggaggttc tgggctgcga ggcacaggac ctctactgac tcgagcctaa 4800  
taaaggaaat ttattttcat tgcaatagtg tggttggttt ttgtgtgcgg ccgcaggaac 4860  
ccctagtgat ggagttggcc actccctctc tgcgcgctcg ctcgctcact gaggccgggc 4920  
gaccaaaggt cgcccgacgc ccgggctttg cccgggcggc ctcagtgagc gagcgagcgc 4980  
gcagctgcct gcaggacat 4999

10

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/24495

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 48/00; C12N 15/63, 15/00; C07H 21/04

US CL : 514/44; 435/320.1, 69.1; 424/93.6; 536/23.51

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/44; 435/320.1, 69.1; 424/93.6; 536/23.51

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN, CAPLUS, MEDLINE, BIOSIS, WEST

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ZATLOUKAL et al. In vivo production of human factor VIII in mice after intrasplenic implantation of primary fibroblasts transfected by receptor-mediated, adenovirus-augmented gene delivery. Proc. Natl. Acad. Sci., U.S.A. May 1994, Vol. 91, No. 11, pages 5148-5152, entire document.	1-17, 20-25, 26, 28
X	CONNELLY et al. Complete short-term correction of canine hemophilia A by in vivo gene therapy. Blood. 15 November 1996, Vol. 88, No. 10, pages 3846-3853, entire document.	1-30
X	CONNELLY et al. Sustained phenotypic correction of murine hemophilia A by in vivo gene therapy. Blood. 01 May 1998, Vol. 91, No. 9, pages 3273-3281, entire document.	1-30

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
*Q* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

27 DECEMBER 1999

Date of mailing of the international search report

10 FEB 2000

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

GAI (JENNIFER) MI LEE

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/24495

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SNYDER et al. Persistent and therapeutic concentrations of human factor IX in mice after hepatic gene transfer of recombinant AAV vector. Nat. Genet. July 1997, Vol. 16, No. 3, pages 270-276, entire document.	1-30
Y	LOZIER et al. Gene therapy and the hemophilias. JAMA, J. Am. Med. Assoc. 04 January 1994, Vol. 271, No. 1, pages 47-51, entire document.	1-30

Form PCT/ISA/210 (continuation of second sheet)(July 1992)★

**THIS PAGE BLANK (USPTO)**



